## **ACITRETIN**

#### **Products Affected**

• acitretin

PA Criteria	Criteria Details
Exclusion Criteria	Severely impaired liver or kidney function. Chronic abnormally elevated blood lipid values. Concomitant use of methotrexate or tetracyclines. Pregnancy. Females of child-bearing potential who intend to become pregnant during therapy or at any time for at least 3 years after discontinuing therapy. Females of child-bearing potential who will not use reliable contraception while undergoing treatment and for at least 3 years following discontinuation. Females of child-bearing potential who drink alcohol during treatment or for two months after cessation of therapy
Required Medical Information	Diagnosis for severe, recalcitrant psoriasis (including plaque, guttate, erythrodermic palmar- plantar and pustular) AND patient must have tried and failed, contraindication or intolerance to one formulary first line agent (e.g., Topical Corticosteroids (betamethasone, fluocinonide, desoximetasone), Topical Calcipotriene/Calcitriol, Topical Calcipotriene, OR Topical Tazarotene)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ACTIMMUNE**

### **Products Affected**

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic granulomatous disease for use in reducing the frequency and severity of serious infections associated with chronic granulomatous disease, or B.) Severe, malignant osteopetrosis (SMO)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ADEMPAS**

#### **Products Affected**

ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form, B.) Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline), C.) Pregnancy, or D.) Patients with pulmonary hypertension associated with idiopathic interstitial pneumonia
Required Medical Information	Diagnosis of one of the following A.) Pulmonary arterial hypertension (WHO group I) and diagnosis was confirmed by right heart catheterization, or B.) Chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) and patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable (Female patients must be enrolled in the ADEMPAS REMS program)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **AFINITOR**

#### **Products Affected**

• AFINITOR DISPERZ

- everolimus
- AFINITOR ORAL TABLET 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Everolimus (Afinitor): Diagnosis of one of the following A.) Advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with Sutent or Nexavar, B.) Diagnosis of pancreatic neuroendocrine tumors (pNET) that are unresectable, locally advanced, or metastatic, C.) Diagnosis of renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery, D.) Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with Femara or Arimidex and the medication will be used in combination with Aromasin, E.) Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection, or F.) Diagnosis of adult patients with progressive, well-differentiated, nonfunctional, neuroendocrine tumors (NET) of gastrointestinal (GI) or lung origin with unresectable, locally advanced, or metastatic disease. Afinitor Disperz: Diagnosis of one of the following A.) Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection, or B.) Diagnosis of partial-onset seizures associated with tuberous sclerosis complex (TSC)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ALECENSA**

### **Products Affected**

ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic anaplastic lymphoma kinase positive non-small cell lung cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ALOSETRON**

## **Products Affected**

• alosetron hcl

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Constipation, B.) History of Chronic or severe constipation or sequelae from constipation, C.) History of ischemic colitis, intestinal obstruction, stricture, toxic megacolon, GI perforation, adhesions, diverticulitis, Crohns disease, ulcerative colitis, D.) History of severe hepatic impairment, E.) History of impaired intestinal circulation, thrombophlebitis, or hypercoagulable state, F.) Coadministration with fluvoxamine
Required Medical Information	Diagnosis of irritable bowel syndrome, severe diarrhea-predominant
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ALPHA1-PROTEINASE INHIBITOR**

#### **Products Affected**

• PROLASTIN-C INTRAVENOUS SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	Not covered for patients with IgA deficiency
Required Medical Information	Diagnosis of alpha-1 proteinase inhibitor (alpha-1 antitrypsin) deficiency in adult patients with emphysema
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ALUNBRIG**

#### **Products Affected**

• ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, ALK positive non-small cell lung cancer and have progressed or are intolerant to Xalkori (crizotinib)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **AMBRISENTAN**

#### **Products Affected**

ambrisentan

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, or B.) Idiopathic pulmonary fibrosis (IPF), including those with pulmonary hypertension
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) and patient has WHO Group I PAH
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **AMPYRA**

### **Products Affected**

• dalfampridine er

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of seizure. B.) Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute)
Required Medical Information	Diagnosis of multiple sclerosis and patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting dalfampridine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **APOKYN**

#### **Products Affected**

• APOKYN SUBCUTANEOUS SOLUTION CARTRIDGE

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with 5-HT(3) receptor antagonists (eg. ondansetron, granisetron, dolasetron, palonosetron, alosetron etc)
Required Medical Information	Diagnosis of Parkinson's disease (PD) and patient is experiencing acute intermittent hypomobility (defined as off episodes characterized by muscle stiffness, slow movements, or difficulty starting movements)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ARCALYST**

### **Products Affected**

ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cryopyrin-associated periodic syndromes (CAPS), including familial cold autoinflammatory syndrome (FCAS) and Muckle-Wells Syndrome (MWS)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ARIKAYCE**

### **Products Affected**

• ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary Mycobacterium avium complex (MAC) infection and used as part of a combination antibacterial regimen in treatment refractory patients
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **AURYXIA**

### **Products Affected**

AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	Iron overload syndrome (e.g. hemochromatosis)
Required Medical Information	Diagnosis of hyperphosphatemia in patients with chronic kidney disease (CKD) on dialysis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or nephrologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **AUSTEDO**

#### **Products Affected**

• AUSTEDO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Suicidal ideation and/or untreated or inadequately treated depression. B.) Hepatic impairment. C. Taking MAOIs, reserpine, or tetrabenazine
Required Medical Information	Diagnosis of one of the following A.) Chorea associated with Huntington's disease (Huntington's chorea), or B.) Tardive dyskinesia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a psychiatrist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **AYVAKIT**

### **Products Affected**

AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic gastrointestinal stromal tumor, with a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BALVERSA**

#### **Products Affected**

BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic urothelial carcinoma with susceptible FGFR3 or FGFR2 genetic alterations and patient has progressed during or following at least one line of prior platinum-containing chemotherapy, including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BANZEL**

### **Products Affected**

BANZEL

PA Criteria	Criteria Details
Exclusion Criteria	Familial Short QT Syndrome
Required Medical Information	Diagnosis of seizures associated with Lennox-Gastaut syndrome
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BENLYSTA**

#### **Products Affected**

• BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of active, autoantibody-positive, system lupus erythematosus (SLE)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BEXAROTENE**

#### **Products Affected**

bexarotene

#### • TARGRETIN EXTERNAL

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BOSENTAN**

#### **Products Affected**

• bosentan

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Receiving concomitant cyclosporine A or glyburide therapy, B.) Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal, or C.) Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND all of the following: A.) Patient has WHO Group I PAH, B.) Patient has New York Heart Association (NYHA) Functional Class II-IV, and C.) Female patients of reproductive potential must use two forms of reliable contraception
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BOSULIF**

### **Products Affected**

• BOSULIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive (Ph+) CML with resistance, relapse, or inadequate response to prior therapy with one of the following tyrosine kinase inhibitors (TKI): Gleevec [imatinib], Tasigna [nilotinib], Sprycel [dasatinib], or B.) newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (Ph + CML)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BRAFTOVI**

#### **Products Affected**

• BRAFTOVI ORAL CAPSULE 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) unresectable or metastic melanoma with documented BRAF V600E or V600K mutation as detected by a FDA-approved test and used in combination with binimetinib, or B.) metastatic colorectal cancer with documented BRAF V600E mutation as detected by an FDA-approved test and patient has received prior therapy. Must be used in combination with cetuximab.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **BRUKINSA**

## **Products Affected**

BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of A.) Mantle Cell Lymphoma (MCL) and patient has tried one prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CABLIVI**

## **Products Affected**

• CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP) and used in combination with plasma exchange and immunosuppression therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	3 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CABOMETYX**

### **Products Affected**

CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	Patients who have or are at risk for severe hemorrhage and/or patients with a recent history of bleeding or hemoptysis
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, or B.) Advanced hepatocellular carcinoma (HCC) and patient has been previously treated with sorafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **CALQUENCE**

### **Products Affected**

• CALQUENCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) MANTLE CELL LYMPHOMA (MCL) and patient has tried one other therapy, B.) Chronic lymphocytic leukemia, or C.) Small lymphocytic lymphoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CAPRELSA**

### **Products Affected**

CAPRELSA

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of medullary thyroid cancer (MTC), and disease is one of the following A.) unresectable, locally advanced, or B.) metastatic AND one of the following: patient has symptomatic disease or patient has progressive disease.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **CARBAGLU**

#### **Products Affected**

• CARBAGLU

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of of one of the following A.) N-acetyl glutamate synthase (NAGS) deficiency AND patient has acute hyperammonemia, or B.) N-acetyl glutamate synthase (NAGS) deficiency AND patient has chronic hyperammonemia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CAYSTON**

### **Products Affected**

• CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	none
Required Medical Information	Diagnosis of cystic fibrosis is confirmed by appropriate diagnostic or genetic testing AND confirmation of P. aeruginosa in cultures of the airways
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CIMZIA**

#### **Products Affected**

• CIMZIA PREFILLED

• CIMZIA SUBCUTANEOUS KIT 2 X 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Active infection, including tuberculosis, B.) Concurrent therapy with other biologics
Required Medical Information	Diagnosis of one of the following A.) Active ankylosing spondylitis (AS) AND patient has trial of/or intolerance/contraindication to Humira and Enbrel, B.) Moderately to severly active Crohn disease AND patient has trial of/or intolerance/contraindication to Humira, C.) Moderate to severe plaque psoriasis AND patient has trial of/or intolerance/contraindication to Humira and Enbrel, D.) Active psoriatic arthritis AND patient has trial of/or intolerance/contraindication to Humira and Enbrel, or E.) Moderately to severely active rheumatoid arthritis (RA) AND patient has trial of/or intolerance/contraindication to Humira and Enbrel, or F.) Non-radiographic axial spondyloarthritis AND patient has had an indequate response to AT LEAST TWO generic Formulary non-steroidal anti-inflammatory drugs (NSAIDs).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CINRYZE**

#### **Products Affected**

• CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Hereditary angioedema (HAE)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CLOBAZAM**

### **Products Affected**

• clobazam

#### SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of seizures associated with Lennox-Gastaut syndrome
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **CNS STIMULANTS**

### **Products Affected**

• armodafinil

modafinil

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Obstructive sleep apnea (OSA) confirmed by sleep lab evaluation, B.) Narcolepsy confirmed by sleep lab evaluation, or C.) Shift work disorder (SWD)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **COMETRIQ**

### **Products Affected**

- COMETRIQ (100 MG DAILY DOSE)
- COMETRIQ (60 MG DAILY DOSE)
- COMETRIQ (140 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following a.) Gastrointestinal perforation, B.) Fistula, or C.) Severe hemorrhage
Required Medical Information	Diagnosis of Progressive, metastatic medullary thyroid cancer (MTC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **COPIKTRA**

#### **Products Affected**

COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory (with history of 2 prior therapies) of one of the following A) chronic lymphocytic leukemia, B) small lymphocytic lymphoma, or C) follicular lymphoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **CORLANOR**

#### **Products Affected**

CORLANOR

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Decompensated acute heart failure, B.) hypotension (i.e. blood pressure less than 90/50 mmHg), C.) sick sinus syndrome or sinoatrial block or 3rd degree AV block (unless a functioning demand pacemaker is present), D.) bradycardia (i.e., resting heart rate less than 60 bpm prior to treatment), or E.) Severe hepatic impairment (Child-Pugh C)
Required Medical Information	Diagnosis of one of the following A.) stable, symptomatic chronic heart failure with left ventricular ejection fraction 35% or less, who are in sinus rhythm with resting heart rate 70 beats per minute or more and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use, or B.) stable, symptomatic heart failure due to dilated cardiomyopathy in patients who are in sinus rhythm with an elevated heart rate
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **COSENTYX**

#### **Products Affected**

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: Failure, contraindication, or intolerance to Enbrel (etanercept) AND Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Cosentyx therapy. All indications (Initial, reauth): Patient is not receiving Cosentyx in combination with a biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab)]. Patient is not receiving Cosentyx in combination with a Janus kinase inhibitor [eg, Xeljanz (tofacitinib)]. For a diagnosis of PsA or plaque psoriasis, Patient is not receiving Cosentyx in combination with a phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)].
Age Restrictions	None
Prescriber Restrictions	Plaque psoriasis (Initial): Prescribed by or in consultation with a dermatologist. PsA (Initial): Prescribed by or in consultation with a rheumatologist or dermatologist. AS (Initial): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	12 months
Other Criteria	None
Indications	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **COTELLIC**

#### **Products Affected**

• COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic malignant melanoma with BRAF V600E OR V600K mutation, and documentation of combination therapy with vemurafenib (Zelboraf)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **CYSTARAN**

#### **Products Affected**

• CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	Demonstrated cysteamine hypersensitivity or penicillamine hypersensitivity
Required Medical Information	Diagnosis of cystinosis and patient has corneal cystine crystal accumulation
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **DALIRESP**

#### **Products Affected**

DALIRESP

PA Criteria	Criteria Details
Exclusion Criteria	Moderate to severe liver impairment (Child-Pugh B or C)
Required Medical Information	Diagnosis of severe chronic obstructive pulmonary disease (COPD) (defined as FEV1 less than or equal to 50% of predicted and FEV1/forced vital capacity [FVC] less than 0.7) associated with chronic bronchitis and a history of COPD exacerbations
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **DAPTOMYCIN**

#### **Products Affected**

daptomycin

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Staphylococcus aureus bacteremia, B.) complicated skin and skin structure infections, including infections caused by methicillin-resistant Staphylococcus aureus (MRSA), or C.) endocarditis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **DAURISMO**

#### **Products Affected**

DAURISMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of newly diagnosed acute myeloid leukemia (AML) and used in combination with cytarabine in patients 75 years of age or older OR in patients that have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **DEFERASIROX**

#### **Products Affected**

- deferasirox oral tablet 360 mg, 90 mg JADENU ORAL TABLET 180 MG
- deferasirox oral tablet soluble
- JADENU SPRINKLE

	]
PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Creatinine clearance less than 40 mL/min, B.) Poor performance status, C.) Platelet count less than 50 x 10(9)/L, D.) Advanced malignancy, E.) High-risk myelodysplastic syndrome (MDS)
Required Medical Information	Diagnosis of one of the following A.) Chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes who have liver iron concentrations of at least 5 mg Fe/g dry weight AND serum ferritin level greater than 300 mcg/L, or B.) Chronic iron overload due to blood transfusions (transfusion hemosiderosis) as evidenced by transfusion of at least 100 mL/kg packed red blood cells AND serum ferritin level greater than 1000 mcg/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **DICLOFENAC TOPICAL**

## **Products Affected**

• diclofenac sodium transdermal gel 3 %

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Actinic keratosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **DRONABINOL**

#### **Products Affected**

dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Anorexia associated to AIDS, or B.) Chemotherapy-induced nausea and vomiting
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **DUPIXENT**

## **Products Affected**

DUPIXENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe atopic dermatitis and patient has trial and failure, contraindication, or intolerance to two medium to high potency topical corticosteroids (e.g., mometasone, triamcinolone, fluocinolone, betamethasone, etc), or B.) Eosinophilic phenotype or oral corticosteroid- dependent moderate to severe asthma and used as an adjunct treatment, or C.) Chronic rhinosinusitis with nasal polyposis and used as an adjunct treatment
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **EMSAM**

#### **Products Affected**

• EMSAM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use with any of the following: SSRIs, SNRIs, clomipramine, imipramine, meperidine, tramadol, methadone, pentazocine, propoxyphene, dextromethorphan, carbamazepine, B.) Pheochromocytoma
Required Medical Information	Diagnosis of major depressive disorder
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ENBREL**

#### **Products Affected**

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SUBCUTANEOUS SOLUTION RECONSTITUTED
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, or E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy. Screening for latent tuberculosis infection is required prior to initiation of treatment.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ENDARI**

#### **Products Affected**

• ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute sickle cell disease AND patient must have trial history of Hydroxyurea. Otherwise Endari requires documentation of (1) history of inadequate treatment with Hydroxyurea OR (2) history of adverse event with Hydroxyurea OR (3) Hydroxyurea is contraindicated.
Age Restrictions	5 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ENTRESTO**

#### **Products Affected**

• ENTRESTO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of angioedema related to previous ACE inhibitor or ARB therapy, B.) Concomitant use or use within 36 hours of ACE inhibitors, or C.) Concomitant use of aliskiren in patients with diabetes
Required Medical Information	Diagnosis of one of the following A.) Chronic heart failure, NYHA Class II to IV, or B.) Symptomatic heart failure with systemic left venticular systolic dysfunction
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **EPIDIOLEX**

#### **Products Affected**

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Lennox-Gastaut syndrome, or B.) Severe myoclonic epilepsy in infancy (Dravet syndrome)
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ERIVEDGE**

## **Products Affected**

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic basal cell carcinoma, or B.) Diagnosis of locally advanced basal cell carcinoma that has recurred following surgery or when the patient is not a candidate for surgery and radiation
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ERLEADA**

#### **Products Affected**

ERLEADA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Nonmetastatic, castration-resistant prostate cancer, or B.) Metastatic, castration-sensitive prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ERLOTINIB**

#### **Products Affected**

• erlotinib hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) locally advanced, unresectable, or metastatic pancreatic cancer and erlotinib (Tarceva) will be used in combination with gemcitabine, or B.) locally advanced or metastatic nonsmall cell lung cancer with one of the following: 1.) failure with at least one prior chemotherapy regimen, 2.) no evidence of disease progression after four cycles of first-line platinum-based chemotherapy and Tarceva will be used as maintenance treatment, or 3.) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ESBRIET**

#### **Products Affected**

- ESBRIET ORAL CAPSULE
- ESBRIET ORAL TABLET 801 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ESRD THERAPY**

#### **Products Affected**

• EPOGEN INJECTION SOLUTION 10000 UNIT/ML, 2000 UNIT/ML, 2000

UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML

RETACRIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Pretreatment hemoglobin levels of less than 10g/dL. Dose reduction or interruption if hemoglobin exceeds 10 g/dL (CKD not on dialysis-adult, cancer), 11 g/dL (CKD on dialysis), 12 g/dL (pediatric CKD) in addition to supporting statement of diagnosis from physician.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **EVEROLIMUS**

#### **Products Affected**

everolimus

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Prevention of kidney transplant organ rejection, or B.) Prevention of liver transplant organ rejection
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescriber is experienced in immunosuppressive therapy and management of transplant patients
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **FARYDAK**

#### **Products Affected**

• FARYDAK ORAL CAPSULE 10 MG, 20 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Multiple Myeloma (MM) Used in combination with both of the following: Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)].
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **FASENRA**

## **Products Affected**

FASENRA

#### • FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe asthma with an eosinophilic phenotype
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FEBUXOSTAT**

#### **Products Affected**

• febuxostat

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of azathioprine or mercaptopurine
Required Medical Information	Diagnosis of Gout and all of the following 1.) documented inadequate treatment response, adverse event, or contraindication to maximally titrated dose of Allopurinol, and 2.) patients with established cardiovascular disease, prescriber attests that benefit of treatment outweighs the risk of treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FENTANYL ORAL**

#### **Products Affected**

• fentanyl citrate buccal lozenge on a handle

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients
Required Medical Information	Must meet all of the following 1.) Diagnosis of cancer-related breakthrough pain, 2.) Patient is currently receiving/tolerant to around-the-clock opioid therapy for persistent cancer pain, and 3.) Patient and prescriber are enrolled in the TIRF REMS Access Program
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# FENTANYL TD

#### **Products Affected**

fentanyl

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients
Required Medical Information	Must meet all of the following 1.) Patient is opioid tolerant (taking for one week or longer at least 60mg of morphine or equivalent daily), and 2.) Patient has tried two extended release oral opioids or is unable to take extended release oral opioids secondary to allergy, adverse events, swallowing difficulty, or uncontrollable nausea/vomiting
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FERRIPROX**

#### **Products Affected**

• FERRIPROX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of transfusional iron overload due to thalassemia syndromes, 2.) Patient has failed prior chelation therapy, and 3.) Patient has an absolute neutrophil count greater than 1.5 x 10(9)/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FIRDAPSE**

## **Products Affected**

• FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	History of seizures
Required Medical Information	Treatment of Lambert-Eaton myasthenic syndrome
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FIRMAGON**

#### **Products Affected**

• FIRMAGON

• FIRMAGON (240 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FORTEO**

#### **Products Affected**

- FORTEO SUBCUTANEOUS SOLUTION 600 MCG/2.4ML
- FORTEO SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following and trial of generic formulary bisphosphonate A.) Osteoporosis in postmenopausal female patient with high risk for fracture, B.) Primary or hypogonadal osteoporosis in male patient with high risk for fracture, or C.) Osteoporosis due to associated sustained systemic glucocorticoid therapy in patient with high risk for fracture
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: 12 months, max treatment 24 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **FYCOMPA**

#### **Products Affected**

• FYCOMPA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Partial-onset seizures with or without secondary generalization, or B.) Primary generalized tonic-clonic seizure disorder
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **GILENYA**

#### **Products Affected**

• GILENYA ORAL CAPSULE 0.5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure, B.) History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker, C.) Baseline QTC interval greater than or equal to 500 milliseconds, D.) Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol)
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **GILOTRIF**

#### **Products Affected**

• GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test, or B.) Metastatic squamous NSCLC, progressing after platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **GLATIRAMER**

#### **Products Affected**

• glatiramer acetate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disese, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **GLEOSTINE**

#### **Products Affected**

• GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Secondary therapy of Hodgkin's disease in combination with other agents and who have relapsed during or failed to respond to primary therapy, B.) Intracranial tumor primary and metastatic who have already received surgical or radiotherapeutic procedures, C.) Carcinoma of the breast, D.) Colorectal cancer, E.) Lung cancer, F.) Malignant melanoma, G.) Malignant tumor of the thymus, H.) Multiple myeloma, I.) Non-Hodgkin's lymphoma, or J.) Small cell carcinoma of lung AND monitoring of blood counts for evidence of Bone Marrow Suppression (thrombocytopenia or leukopenia).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **GOCOVRI**

#### **Products Affected**

• GOCOVRI

PA Criteria	Criteria Details
Exclusion Criteria	Patients with end-stage reanl disease (ESRD, CrCl below 15 ml/min/m2)
Required Medical Information	Diagnosis of one of the following A.) Parkinsons disease and patient is experiencing dyskinesia, receiving levodopa based therapy, and has documented trial and failure to amantadine immediate release, or B.) Extrapyramidal disease and has documented trial and failure to amantadine immediate release
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **GROWTH HORMONE**

#### **Products Affected**

OMNITROPE

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) growth promotion in pediatric patients with closed epiphyses, B.) acute critical illness caused by complications following open-heart or abdominal surgery, multiple accidental trauma, or acute respiratory failure, C.) active malignancy, D.) active proliferative or servere nonproliferative diabetic retinopathy
Required Medical Information	Diagnosis of pediatric indication: A.) GHD and bone age at least 1 year or 2 standard deviations (SD) delayed compared with chronological age and 2 stim tests with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SD below mean if CNS pathology, h/o irradiation, or proven genetic cause, B.) SGA and birth weight or length 2 or more SD below mean for gestational age and fails to manifest catch up growth by age 2 (height 2 or more SD below mean for age and gender), C.) CRI and nutritional status has been optimized, metabolic abnormalities have been corrected, and patient has not had renal transplant, D.) SHOX deficiency or Noonan syndrome, E.) PWS confirmed by genetic testing, F.) Turner Syndrome confirmed by chromosome analysis. Diagnosis of GHD, CRI, SHOX deficiency, Noonan syndrome, and PWS one of the following: 1.) height more than 3 SD below mean for age and gender, 2.) height more than 2 SD below mean with GV more than 1 SD below mean, or 3.) GV over 1 year 2 SD below mean. Diagnosis of adult indication: A.) childhood or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone, glucagon, arginine), B.) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C.) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D.) GHD with irreversible hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region, a subnormal IGF-1 (after at least 1 month off GH therapy), objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or cardiovascular complications,

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
	completed linear growth (GV less than 2 cm/year), and GH has been discontinued for at least 1 month (if previously receiving GH).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **HEPATITIS B**

#### **Products Affected**

adefovir dipivoxil

BARACLUDE ORAL SOLUTION

- entecavir
- VEMLIDY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic hepatitis B and all of the following: 1.) Patient has evidence of viral replication, 2.) Patient has evidence of persistent elevations in serum aminotransferase (ALT or AST) or histologically active disease, and 3.) Patient is receiving anti-retroviral therapy if the patient has HIV co-infection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **HEPATITIS C**

#### **Products Affected**

MAVYRET

• VOSEVI

• sofosbuvir-velpatasvir

- Sofese with temperature in the second seco	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of HCV genotype, subtype and quantitative HCV RNA (viral load) testing any time prior to therapy. Must document the following within 12 weeks of starting therapy: CBC, INR, hepatic function panel, and GFR. Must document cirrhosis status, prior treatment history (if any), and planned duration of treatment. All genotypes will require trial/failure, contraindication to, or intolerance to Mavyret or Sofosbuvir-Velpatasvir prior to the approval of Vosevi.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in conjunction with gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **HUMIRA**

#### **Products Affected**

- **HUMIRA PEDIATRIC CROHNS** START SUBCUTANEOUS PREFILLED • HUMIRA PEN-PS/UV/ADOL HS SYRINGE KIT 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML
- HUMIRA PEN SUBCUTANEOUS PEN-**INJECTOR KIT**
- HUMIRA PEN-CD/UC/HS STARTER
- **START**
- HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT

INJECTOR KIT	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy, G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants (e.g. corticosteroids, azathioprine), H.) Noninfectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa. Screening for latent tuberculosis infection is required prior to initiation of treatment.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135 PA20 C

Formulary ID:20153 Version 12

Y0135\_PA20\_C Formulary ID:20153\_Version 12

# **IBRANCE**

#### **Products Affected**

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy, or B) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with an aromatase inhibitor AND One of the following 1) patient is a postmenopausal woman, 3) patient is a man, or 3) both of the following: patient is a premenopausal or perimenopausal woman and patient is receiving a luteinizing hormone-releasing hormone (LHRH) agonist [eg, Zoladex (goserelin)].
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ICLUSIG**

#### **Products Affected**

• ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated, or B.) Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **IDHIFA**

#### **Products Affected**

• IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia with an isocitrate dehydrogenase 2 mutation as detected by an FDA approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **IMATINIB**

#### **Products Affected**

• imatinib mesylate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), B.) Ph+ acute lymphoblastic leukemia (ALL), C.) Gastrointestinal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, D.) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, E.) hypereosinophilic syndrome or chronic eosinophilic leukemia, F.) myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, or G.) aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **IMBRUVICA**

### **Products Affected**

• IMBRUVICA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) who have received at least one prior therapy, B.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL), C.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion, D.) Waldenstrom's macroglobulinemia (WM), E.) Marginal zone lymphoma who require systemic therapy and have received at least one prior anti-CD20-based therapy, or F.) Graft vs host disease after failure of a least one first-line corticosteroid therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **INCRELEX**

### **Products Affected**

• INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) active or suspected malignancy, B.) use for growth promotion in patients with closed epiphyses, C.) Intravenous administration
Required Medical Information	Diagnosis of one of the following A.) growth failure in children with severe primary IGF-1 deficiency, or B.) growth hormone (GH) gene deletion in children who have developed neutralizing antibodies to GH
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **INHALED TOBRAMYCIN**

#### **Products Affected**

• tobramycin inhalation

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of cystic fibrosis and 2.) Patient has evidence of P. aeruginosa in the lungs
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **INLYTA**

#### **Products Affected**

• INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced renal cell carcinoma and patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **INREBIC**

#### **Products Affected**

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate-2 or high-risk primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis (MF).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **INTRAROSA**

#### **Products Affected**

• INTRAROSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, B.) Known or suspected estrogendependent neoplasia
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe dyspareunia due to menopause, or B.) Atrophic vaginitis due to menopause.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 3 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **INTRON A**

#### **Products Affected**

• INTRON A

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis, or B.) Decompensated liver disease
Required Medical Information	Diagnosis of one of the following A.) Hairy cell leukemia, B.) Diagnosis of condylomata acuminata involving external surfaces to the genital or perianal areas, C.) Diagnosis of AIDS-related Kaposi's sarcoma, D.) Clinically aggressive follicular lymphoma and the medication will be used concurrently with anthracycline-containing chemotherapy or is not a candidate for anthracycline-containing chemotherapy, E.)Malignant melanoma and the request for coverage is within 56 days of surgery and the patient is at high risk of disease recurrence, F.) Diagnosis of chronic hepatitis B with compensated liver disease and patient has evidence of hepatitis B viral replication and patient has been serum hepatitis B surface antigen-positive for at least 6 months, or G.) Diagnosis of chronic hepatitis C with compensated liver disease and is receiving combination therapy with ribavirin, unless ribavirin is contraindicated, and the medication will not be used as part of triple therapy with a protease inhibitor and patient has a clinical reason for not using peginterferon
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Condylomata: 3 mos, HBV: E antigen pos: 16 wks, E antigen neg: 48 wks, KS: 16 wks, Other: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **IRESSA**

#### **Products Affected**

• IRESSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic Non-small cell lung cancer (NSCLC) and Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ITRACONAZOLE**

#### **Products Affected**

• itraconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.)
Required Medical Information	Diagnosis of one of the following A.) Systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), B.) Onychomycosis confirmed by one of the following: positive potassium hydroxide (KOH) preparation, fungal culture, or nail biopsy, or C.) Candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole (ORAL SOLUTION ONLY)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **IVIG**

#### **Products Affected**

- GAMMAGARD INJECTION SOLUTION 2.5 GM/25ML
- GAMMAGARD S/D LESS IGA
- GAMMAKED INJECTION SOLUTION 1 GM/10ML
- GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML
- GAMUNEX-C INJECTION SOLUTION 1 GM/10ML
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 2 GM/20ML

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Acute corn or maltose hypersensitivity, B.) Hereditary fructose intolerance, C.) Hyperprolinemia, D.) IgA deficiency with antibody formation and a history of hypersensitivity, E.) History of anaphylaxis or severe systemic reaction to human immune globulin
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153 Version 12

# **JAKAFI**

#### **Products Affected**

JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Intermediate or high-risk myelofibrosis, or B.) Polycythemia vera: Diagnosis of polycythemia vera, AND history of failure, contraindication, or intolerance to hydroxyurea, or C.) Acute Graft Versus Host Disease (GVHD): Diagnosis of Acute GVHD, AND disease is refractory to steroid therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **JYNARQUE**

### **Products Affected**

• JYNARQUE ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) use in patients unable to sense or respond to thirst, B.) anuria, C.) history, signs, or symptoms of significant liver impairment or injury, D.) uncorrected abnormal blood sodium concentrations, E.) uncorrected urinary outflow obstruction
Required Medical Information	Diagnosis of autosomal dominant polycystic kidney disease (ADPKD)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **KALYDECO**

### **Products Affected**

KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis AND the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **KISQALI**

### **Products Affected**

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)
- KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KISQALI FEMARA(200 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	KISQALI: Breast Cancer: Diagnosis of one of the following A.) Metastatic or advanced, HER-2 negative, hormone receptor-positive, postmenopausal women in combination with fulvestrant as initial endocrine based therapy or following disease progression on endocrine therapy, or B.) Metastatic or advanced, HER-2 negative, hormone receptor-positive, premenopausal, perimenopausal or postmenopausal women, in combination with an aromatase inhibitor for initial endocrine-based treatment. KISQALI FEMARA: Diagnosis of HER-2 negative, hormone receptor-positive, advanced or metastatic breast cancer in premenopausal, perimenopausal, or postmenopausal women, as initial endocrine based therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **KORLYM**

### **Products Affected**

• KORLYM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, C.) concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, D.) history of unexplained vaginal bleeding, E.) endometrial hyperplasia with atypia or endometrial carcinoma
Required Medical Information	Diagnosis of endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and one of the following A.) Used to control hyperglycemia secondary to hypercortisolism and patient has failed surgery, or B.) Used to control hyperglycemia secondary to hypercortisolium and patient is not a candidate for surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **KUVAN**

#### **Products Affected**

• KUVAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **LENVIMA**

#### **Products Affected**

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer, B.) Advanced renal cell carcinoma following one prior anti-angiogenic therapy in combination with everolimus, C.) Unresectable liver carcinoma, or D.) Advanced endometrial carcinoma that is not microsatellite instability-high or mismatch repair deficient, in a patient which has disease progression following prior systemic therapy and is not a candidate for curative surgery or radiation
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135 PA20 C

Formulary ID:20153 Version 12

# **LEUKINE**

#### **Products Affected**

• LEUKINE INJECTION SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with myelosuppresive chemotherapy or radiation or excessive (greater than or equal to 10%) leukemic myeloid blasts in bone marrow or peripheral blood
Required Medical Information	Diagnosis of one of the following A.) Patient has undergone allogeneic or autologous bone marrow transplant (BMT) and engraftment is delayed or failed, B.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, C.) Medication will be used for myeloid reconstitution after an autologous or allogeneic BMT, D.) Patient has acute myeloid leukemia and administration will be after completion of induction chemotherapy, or E.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **LEUPROLIDE**

#### **Products Affected**

- ELIGARD
- leuprolide acetate injection
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) advanced or metastatic prostate cancer and patient with have trial of/contraindication to Eligard prior to approval of Lupron, B.) Diagnosis of central precocious puberty and patient had early onset of secondary sexual characteristics (male: earlier than 9 years of age. female: earlier than 8 years of age) and advanced bone age of at least one year compared with chronological age and has undergone gonadotropin-releasing hormone agonist (GnRHa) testing with peak luteinizing hormone (LH) level above pre-pubertal range or random LH level in pubertal range and Patient had the following diagnostic evaluations to rule out tumors, when suspected: diagnostic imaging of the brain (MRI or CT scan), Pelvic/testicular/adrenal ultrasound, Human chorionic gonadotropin levels, Adrenal steroids to rule out congenital adrenal hyperplasia, C.) the medication will be used for stimulation testing to confirm the diagnosis of central precocious puberty, D.) management of endometriosis, or E.) anemia caused by uterina leiomyomata
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **LIDOCAINE EXT**

#### **Products Affected**

- lidocaine external ointment
- lidocaine hcl external solution
- lidocaine hcl urethral/mucosal external gel
- lidocaine-prilocaine external cream

PA Criteria	Criteria Details
Exclusion Criteria	Amide hypersensitivity
Required Medical Information	For topical anesthesia of skin and mucous membranes
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# LIDOCAINE PATCH

#### **Products Affected**

• lidocaine external patch 5 %

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) pain associated with diabetic neuropathy, B.) pain associated with cancer-related neuropathy, C.) post-herpetic neuralgia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **LINEZOLID**

#### **Products Affected**

- linezolid intravenous solution 600 mg/300ml
- linezolid oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Linezolid should not be used concurrently or within 14 days of MAOI therapy.
Required Medical Information	Supporting statement of diagnosis from the physician OR susceptibility testing shows drug activity for infection being treated
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	VRE: 4 weeks. Nosocomial and community acquired pneumonia: 3 weeks. All other indications: 2 weeks
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **LONSURF**

#### **Products Affected**

• LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy, or B.) Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least 2 prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy if appropriate
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **LORBRENA**

#### **Products Affected**

LORBRENA

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A4 inducers
Required Medical Information	Diagnosis of metastatic, anaplastic lymphoma kinase (ALK) positive non-small cell lung cancer with disease progression on either alectinib or ceritinib as the first ALK inhibitor for metastatic disease, or disease progression on crizotinib and at least one other ALK inhibitor for metastatic disease
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **LYNPARZA**

#### **Products Affected**

• LYNPARZA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) HER2- negative, deleterious or suspected deleterious germline BRCA mutated metastatic breast cancer AND patient has been previously treated with chemotherapy in neoadjuvant, adjuvant, or metastatic setting, B.) Advanced ovarian cancer with known or suspected BRCA mutation as detected by an FDA-approved test AND patient has trial and failure, contraindication, or intolerance to 3 or more prior lines of chemotherapy, C.) Recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer AND used for maintenance treatment in patients who are in complete or partial response to platinum-based chemotherapy (e.g. cisplatin, carboplatin), D.) Deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients with complete or partial response to first-line platinum-based chemotherapy, or E.) Deleterious or suspected deleterious germline BRCA-mutated metastatic pancreatic adenocarcinoma and disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **MATULANE**

#### **Products Affected**

• MATULANE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Hodgkin's Disease, Stages III and IV in combination with other anticancer drugs, B.)Malignant intracranial tumor including but not limited to medulloblastoma, C.) Multiple myeloma, D.) Non-Hodgkin's lymphona, or E.) Malignant glioma
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **MAYZENT**

#### **Products Affected**

MAYZENT

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) CYP2C9*3/*3 genotype, B.) In the last 6 months experienced myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, Class III-IV heart failure, or C.) Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome, unless patient has a functioning pacemaker
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis, including clinically isolated syndrome, relapsing-remitting disease, or active secondary progressive disease and the following A.) Patients with relapsing forms of multiple sclerosis have history of/or contraindication to Avonex, Betaseron, Copaxone, or Gilenya
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **MEKINIST**

#### **Products Affected**

MEKINIST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and used in combination with dabrafenib and no locoregional treatment options, B.) Malignant melanoma with lymph node involvement and following complete resection with BRAF V600E or V600K mutations and used in combination with dabrafenib, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutations and used in combination with dabrafenib or as monotherapy, or D.) Metastatic nonsmall cell lung cancer, with BRAF V600E mutation, in combination with dabrafenib.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **MEKTOVI**

#### **Products Affected**

MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic malignant melonoma with documented BRAF V600E or V600K mutation as detected by an FDA approved test AND used in combination with encorafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

### **METHOTREXATE SC**

#### **Products Affected**

- OTREXUP SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.4ML, 12.5 MG/0.4ML, 15 MG/0.4ML, 17.5 MG/0.4ML, 20 MG/0.4ML, 22.5 MG/0.4ML, 25 MG/0.4ML
- RASUVO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.2ML, 12.5 MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML

PA Criteria	Criteria Details
Exclusion Criteria	A.) Pregnancy, B.) Breastfeeding, C.) Alcoholism or liver disease, D.) Immunodeficiency syndromes, E.) Preexisting blood dyscrasias
Required Medical Information	Diagnosis of one of the following: A.) Severe rheumatoid arthritis in patients who are intolerant of or had an inadequate response to first-line therapy, B.) Polyarticular juvenile idiopathic arthritis in patients who are intolerant of or had an inadequate response to first-line therapy, C.) Severe, recalcitrant, disabling psoriasis in patients who are not adequately responsive to other forms of therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **METHOXSALEN**

#### **Products Affected**

• methoxsalen rapid

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Aphakia, B.) Melanoma or a history of melanoma, C.) Invasive squamous cell carcinomas, or D.) History of a light sensitive disease/skin photosensitivity disorder such systemic lupus erythematosus (SLE), porphyria cutanea tarda, erythropoietic protoporphyria, variegate porphyria, xeroderma pigmentosum or albinism
Required Medical Information	Diagnosis of one of the following A.) Psoriasis, B.) Cutaneous T-cell lymphoma, or C.) Vitiligo
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, immunologist, or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **MIGLUSTAT**

#### **Products Affected**

• miglustat

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **MS INTERFERONS**

#### **Products Affected**

- AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT
- AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **MYTESI**

#### **Products Affected**

• MYTESI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of non-infectious diarrhea associated with HIV/AIDS in patients receiving anti-retroviral therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or gastroenterologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NATPARA**

#### **Products Affected**

NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hypoparathyroidism and used to control hypocalcemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NERLYNX**

#### **Products Affected**

NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) early stage HER2-positive breast cancer and used after trastuzumab therapy, or B.) advanced or metastatic HER2-positive breast cancer and patient has received 2 or more prior anti-HER2-based regimens in the metastatic setting
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **NEXAVAR**

#### **Products Affected**

NEXAVAR

PA Criteria	Criteria Details
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Diagnosis of locally recurrent or metastatic, progressive, differentiated thyroid carcinoma that is refractory to radioactive iodine treatment, or C.) Diagnosis of unresectable hepatocellular carcinoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NINLARO**

#### **Products Affected**

NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma and documentation of combination therapy with lenalidomide and dexamethasone, used in patients with history of 1 prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NORTHERA**

#### **Products Affected**

NORTHERA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (e.g., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NOXAFIL**

#### **Products Affected**

• NOXAFIL ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids
Required Medical Information	Diagnosis of one of the following A.) Oropharyngeal candidiasis, or B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection
Age Restrictions	13 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NUBEQA**

#### **Products Affected**

NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of non-metastatic, castration-resistant prostate cancer
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NUCALA**

#### **Products Affected**

NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Severe asthma with eosinophilic phenotype, or B.) Eosinophilic granulomatosis with polyangiitis (EGPA)
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist, rheumatologist, or immunologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **NUEDEXTA**

#### **Products Affected**

NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of prolonged QT interval, congenital long QT syndrome or Torsades de pointes, B.) Heart failure, C.) Complete AV block without an implanted pacemaker or high risk of complete AV block, D.) Concomitant use with quinidine, quinine, mefloquine, or drugs that prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), E.) Concomitant use with MAOIs or within 14 days of MAOI therapy
Required Medical Information	Diagnosis of pseudobulbar affect
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **OCTREOTIDE**

#### **Products Affected**

• octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) acromegaly and patient has inadequate response to or is ineligible for surgery, radiation, or bromocriptine mesylate, B.) metastatic carcinoid syndrome, C.) vasoactive intestinal peptide-secreting tumors (VIPomas) with associated diarrhea
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ODOMZO**

#### **Products Affected**

• ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of locally advanced basal cell carcinoma of the skin and one of the following A.) Cancer has recurred following surgery or radiation therapy, B.) Patient is not a candidate for surgery or radiation therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **OPSUMIT**

#### **Products Affected**

• OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension WHO group I AND diagnosis was confirmed by right heart catheterization AND female patients are enrolled in the OPSUMIT REMS program
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ORFADIN**

#### **Products Affected**

ORFADIN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of tyrosinemia type 1
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ORKAMBI**

#### **Products Affected**

ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **OSPHENA**

### **Products Affected**

OSPHENA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) undiagnosed abnormal genital bleeding, B.) known or suspected estrogen-dependent neoplasia, C.) active or history of DVT, D.) active or history of pulmonary embolism, E.) active or history of arterial thromboembolic disease F.) pregnancy
Required Medical Information	Diagnosis of one of the following A.) moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause, or B.) moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **OXANDROLONE**

#### **Products Affected**

• oxandrolone oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Breast or prostate cancer in men, B.) Breast cancer in women with hypercalcemia, C.) Pregnancy, D.) Nephrosis or nephrotic phase of nephritis, E.) Hypercalcemia
Required Medical Information	Diagnosis one of the following and receiving treatment as an adjunct therapy to promote weight gain A.) Extensive surgery, B.) Chronic infections, C.) Severe trauma, or D.) Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons, E.) Chronic corticosteroid administration, F.) Bone pain associated with osteoporosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **PANRETIN**

### **Products Affected**

PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Kaposi sarcoma cutaneous lesions in patient with AIDS-related Kaposi sarcoma (KS)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or HIV specialist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **PCSK9 INHIBITOR**

#### **Products Affected**

- PRALUENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REPATHA

- REPATHA PUSHTRONEX SYSTEM
- REPATHA SURECLICK

• KEFATHA	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	PRALUENT: Must meet criteria #1, #2 or #3. REPATHA: Must meet criteria #1, #2, #3 or #4. 1.) Diagnosis of primary hyperlipidemia including heterozygous familial hypercholesterolemia (HeFH). 2.) Myocardial infarction prophylaxis, stroke prophylaxis, and to reduce risk of coronary revascularization in pts with established CVD. 3.) Diagnosis of clinical atherosclerotic cardiovascular disease (CVD) as defined as one of the following: a. acute coronary syndrome, b. history of myocardial infarction, c. stable/unstable angina, d. coronary or other arterial revascularization, e. stroke, f. transient ischemic stroke (TIA), g. peripheral arterial disease presumed to be atherosclerotic region. 4.) Primary hyperlipidemia homozygous familial hypercholesterolemia (HoFH) confirmed by genotyping OR diagnosis based on the following: a. History of untreated LDL-C greater than 500 mg/dL AND xanthoma before 10 years of age OR b. Documentation of HeFH in both parents. REQUIRED DOCUMENTATION FOR INITIAL THERAPY: A.) Baseline and current LDL-C, LDL-C greater than or equal to 70 mg/dL, AND used in combination with maximally tolerated high-intensity statin OR patient is statin intolerant and LDL-C greater than or equal to 70 mg/dL. FOR CONTINUING THERAPY: Will continue to be used in combination with maximally tolerated statin (unless statin intolerant).
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist
Coverage Duration	Initial: 8 weeks, Renewal: 12 months

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **PEGYLATED INTERFERON**

#### **Products Affected**

- PEGASYS PROCLICK SUBCUTANEOUS SOLUTION 180 MCG/0.5ML
- PEGASYS SUBCUTANEOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon, B.) Uncontrolled depression
Required Medical Information	Diagnosis of one of the following A.) Chronic hepatitis B infection, or B.) Chronic hepatitis C and required criteria will be applied consistent with current AASLD-IDSA guidance with compensated liver disease
Age Restrictions	Hepatitis B: 3 years of age and older. Hepatitis C: 5 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist, gastroenterologist, or infectious disease specialist
Coverage Duration	HBV: 12 months, HCV: based on current AASLD guidelines
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **PENICILLAMINE**

### **Products Affected**

• penicillamine oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Breastfeeding, B.) During Pregnancy (except for treatment of Wilson's disease), C.) Hypersensitivity to penicillamine products, D.) Penicillamine-related aplastic anemia/agranulocytosis, E.) Rheumatoid arthritis patients with history or evidence of renal insufficiency
Required Medical Information	Diagnosis of one of the following A.) Cystinuria, B.) Rheumatoid arthritis,or C.) Wilson's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **PIQRAY**

#### **Products Affected**

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)

• PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR) positive, HER2-negative, PIK3CA-mutated, advanced or metastatic breast cancer and used in combination with fulvestrant for postmenopausal women, and men following progression on or after endocrine- based regimen.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **POMALYST**

#### **Products Affected**

POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Must meet all of the following 1.) Disease has progressed on or within 60 days of completion of the last therapy, 2.) If female of reproductive potential ALL of the below: Two negative pregnancy tests obtained prior to initiating therapy with Pomalyst, monthly negative pregnancy tests during therapy, 3.) Patient has been counseled about the use of 2 forms of reliable contraception before, during, and 1 month after discontinuing therapy with Pomalyst, 4.) Patient assessment to determine if prophylactic aspirin or antithrombic treatment (warfarin, clopidogrel) will need to be taken to reduce the risk of VTE (embolism, stroke), and 5.) Registered and certified to be compliant with Pomalyst REMS program
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **POSACONAZOLE**

#### **Products Affected**

posaconazole

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids
Required Medical Information	Diagnosis of one of the following A.) Oropharyngeal candidiasis, or B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection
Age Restrictions	13 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **PROMACTA**

#### **Products Affected**

• PROMACTA ORAL PACKET 12.5 MG • PROMACTA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic thrombocytopenic purpura (ITP), B.) Chronic hepatitis C infection associated thrombocytopenia, or C.) Severe aplastic anemia with insufficient response to immunosuppressive therapy or in combination with standard immunosuppressive therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **PULMONARY FIBROSIS**

#### **Products Affected**

OFEV

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Idiopathic pulmonary fibrosis (IPF), B.) Systemic sclerosis-associated interstitial lung disease (ILD), or C.) Chronic fibrosing interstitial lung disease with a progressive phenotype
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **PURIXAN**

#### **Products Affected**

• PURIXAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute lymphoblastic leukemia AND failure of mercaptopurine tablets unless contraindicated or clinically significant adverse effects are experienced, AND OR member has a documented swallowing disorder or an inability to swallow tablets or capsules.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **QUININE SULFATE**

#### **Products Affected**

• quinine sulfate oral

PA Criteria	Criteria Details
Exclusion Criteria	Prolongation of QT interval. Glucose-6-phosphate dehydrogenase deficiency. Myasthenia gravis. Known hypersensitivity to mefloquine or quinidine. Optic neuritis. Diagnosis of Blackwater fever
Required Medical Information	Diagnosis of one of the following A.) uncomplicated Plasmodium falciparum malaria, B.) uncomplicated Plasmodium vivax malaria, or C.) babesiosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **RANOLAZINE**

#### **Products Affected**

• ranolazine er

PA Criteria	Criteria Details
Exclusion Criteria	A.) Hepatic cirrhosis, B.) Concurrent therapy with a strong CYP3A4 inhibitor, C.) Concurrent therapy with a CYP3A4 inducer
Required Medical Information	Diagnosis of chronic angina
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **RAVICTI**

#### **Products Affected**

• RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of urea cycle disorders
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **REGRANEX**

#### **Products Affected**

• REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	Known neoplasm at the site of application
Required Medical Information	Diagnosis of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **REVLIMID**

#### **Products Affected**

• REVLIMID

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality or without additional cytogenetic abnormalities, D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **RILUTEK**

#### **Products Affected**

• riluzole

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of amyotrophic lateral sclerosis (ALS)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **RINVOQ**

#### **Products Affected**

• RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of moderate to severe rheumatoid arthritis and patient has had an inadequate response or intolerance to methotrexate
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ROZLYTREK**

## **Products Affected**

ROZLYTREK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) ROS1-positive metastatic non-small cell lung cancer (NSCLC), OR B) Solid tumors that 1) have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, AND 2) are metastatic or where surgical resection is likely to result in severe morbidity, AND 3) have either progressed following treatment or have no satisfactory alternative therapy
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **RUBRACA**

#### **Products Affected**

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of 1. deleterious BRCA mutation (germline and/or somatic)-associated ovarian, fallopian tube, or primary peritoneal cancer and all of the following criteria (A-E): A.) BRCA mutation positive as detected by an approved FDA laboratory test, B.) Previous trial/failure with two or more chemotherapy regimens, C.) Used as monotherapy, D.) Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, E.) Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose. Diagnosis of 2. Diagnosis of recurrent ovarian, fallopian tube, or primary peritoneal cancer and all of the following (A-D): A.) Complete or partial response to platinum-based chemotherapy B.) Used as monotherapy C.) Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, D.) Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose. Renewal will be based on lack of disease progression or unacceptable toxicity.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

Y0135\_PA20\_C Formulary ID:20153\_Version 12 Last Updated: 05/19/2020

Effective Date: 06/01/2020

# **RUCONEST**

#### **Products Affected**

• RUCONEST

PA Criteria	Criteria Details
Exclusion Criteria	Known allergy to rabbits or rabbit-derived products (leporine protein hypersensitivity)
Required Medical Information	Diagnosis of Hereditary angioedema (HAE)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **RYDAPT**

#### **Products Affected**

• RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and consolidation therapy, or B.) systemic mastocytosis or mast cell leukemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SAMSCA**

#### **Products Affected**

• SAMSCA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) use in patients unable to sense or respond to thirst, B.) anuria, C.) hypovolemic hyponatremia, D.) urgent need to raise serum sodium acutely
Required Medical Information	Diagnosis of clinically significant hypervolemic and euvolemic hyponatremia (serum sodium less than 125 mEq/L or less marks hyponatremia that is symptomatic and has resisted correction with fluid restriction), including in patients with heart failure and syndrome of inappropriate antidiuretic hormone (SIADH)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SIGNIFOR**

#### **Products Affected**

SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing disease and patient has inadequate response to or is not a candidate for surgery
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **SILDENAFIL**

#### **Products Affected**

• sildenafil citrate oral tablet 20 mg

PA Criteria	Criteria Details
Exclusion Criteria	Nitrate therapy
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) and Patient has WHO Group I PAH
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SKYRIZI**

#### **Products Affected**

• SKYRIZI (150 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of moderate to severe plaque psoriasis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SOLTAMOX**

#### **Products Affected**

• SOLTAMOX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant coumarin-type anticoagulant therapy, B.) history of thromboembolic disease such as DVT or PE
Required Medical Information	Diagnosis of breast cancer and documentation of inability to swallow tablet formulation
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SOMATULINE**

#### **Products Affected**

• SOMATULINE DEPOT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) acromegaly in patient with inadequate response to or is ineligible for surgery or radiotherapy, B.) carcinoid syndrome, or C.) gastroenteropancreatic neuroendocrine tumors (GEP-NETs)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SOMAVERT**

#### **Products Affected**

• SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly and patient has inadequate response to or is ineligible for surgery or radiation therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SPRYCEL**

#### **Products Affected**

SPRYCEL

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) that is newly diagnosed in the chronic phase, B.) Ph+ CML in chronic, accelerated, or lymphoid blast phase with resistance or intolerance to prior therapy, C.) Diagnosis of Ph+ acute lymphoblastic leukemia with resistance or intolerance to prior therapy, or D.) Newly diagnosed Ph+ acute lymphoblastic leukemia in combination with chemotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **STELARA**

#### **Products Affected**

- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) moderate to severely active Crohn disease and patient has trial and failure or intolerance or contraindication to Humira, B.) moderate to severe plaque psoriasis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, C.) active psoriatic arthritis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, or D.) moderate to severe active ulcerative colitis and patient has trial and failure or intolerance or contraindication to Humira
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist or gastroenterologist or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **STIVARGA**

#### **Products Affected**

STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) metastatic colorectal cancer in patients previously treated with ALL of the following per the indication: 1. (fluoropyrimidine-, oxaliplatin-, and irinotecan)-based chemotherapy 2. anti-VEGF bevacizumab 3. anti-EGFR panitumumab OR cetuximab (for KRAS mutation-negative patients only), B.) liver carcinoma in patients previously treated with sorafenib, or C.) locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) who have been previously treated with imatinib and sunitinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SUTENT**

## **Products Affected**

• SUTENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) gastrointestinal stromal tumor after disease progression on or intolerance to imatinib, B.) pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease, C.) advanced renal cell carcinoma, or D.) renal cell carinoma and used as adjuvant therapy following nephrectomy in patients who are at high risk for recurrence
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SYLATRON**

#### **Products Affected**

• SYLATRON SUBCUTANEOUS KIT 200 MCG, 300 MCG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis or B.) Hepatic decompensation (Child-Pugh score greater than 6 [Class B or C])
Required Medical Information	Diagnosis of melanoma with microscopic or gross nodal involvement and prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SYMDEKO**

#### **Products Affected**

SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis and One of the following A.) Patient is homozygous for the F508del mutation, or B.) Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **SYMLIN**

#### **Products Affected**

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) confirmed diagnosis of gastroparesis, B.) hypoglycemia unawareness
Required Medical Information	Diagnosis of one of the following A.) type 1 diabetes mellitus and patient uses mealtime insulin therapy and has failed to achieve desired glucose control, or B.) type 2 diabetes mellitus and patient uses mealtime insulin therapy and has failed to achieve desired glucose control
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SYNAREL**

## **Products Affected**

SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) breastfeeding, C.) undiagnosed abnormal vaginal bleeding
Required Medical Information	Diagnosis of one of the following A.) central precocious puberty, or B.) endometriosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SYNRIBO**

## **Products Affected**

SYNRIBO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic or accelerated phase chronic myeloid leukemia (CML) and patient has tried and failed or has a contraindication or intolerance to 2 tyrosine kinase inhibitors
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **SYPRINE**

#### **Products Affected**

CLOVIQUE

• trientine hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease in patients that are intolerant to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TAFINLAR**

#### **Products Affected**

TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	A.) Diagnosis of locally advanced or metastatic anaplastic thyroid carcinoma with BRAF V600E mutation, in combination with trametinib and no satisfactory locoregional treatment options OR B.) Diagnosis of metastatic non-small cell lung cancer with BRAF V600E mutation, in combination with trametinib OR in patients previously treated as monotherapy OR C.) Diagnosis of unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutation AND 1) used as monotherapy OR 2) in combination with trametinib OR 3) used as adjuvant therapy following complete resection in patients with lymph node involvement AND used in combination with trametinib.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TAGRISSO**

#### **Products Affected**

• TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) metastatic, non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R mutation and used as first line therapy, or B.) Metastatic, non-small cell lung cancer with confirmed presence of T790M EGFR mutation AND whose disease has progressed on or after EGFR tyrosine kinase inhibitor based therapy (Diagnosis should be confirmed by an FDA-approved test)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TALZENNA**

#### **Products Affected**

TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious or suspected deleterious germline BRCA-mutated (gBRCAm), HER2-negative locally advanced or metastatic breast cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TASIGNA**

#### **Products Affected**

TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia, or D.) Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed Philadelphia chromosome-positive chronic myelogenous leukemia (CML) in chronic phase, B.) Chronic-phase and accelerated-phase Philadelphia chromosome-positive CML in patients resistant or intolerant to prior therapy that include imatinib, or C.) Chonic phase Phildelphia chromosome-positive CML in patients with resistance or intolerance to prior tyrosine-kinase inhibitor therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TAVALISSE**

### **Products Affected**

• TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of thrombocytopenia in patient with chronic idiopathic thrombocytopenic purpura (ITP) and an insufficient response to one previous treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TAZORAC**

#### **Products Affected**

• tazarotene external

- TAZORAC EXTERNAL GEL
- TAZORAC EXTERNAL CREAM 0.05 %

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) acne vulgaris and patient has trial with at least one generic topical acne product, or B.) stable moderate to severe plaque psoriasis with 20% or less body surface area involvement and patient has trial with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TAZVERIK**

#### **Products Affected**

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or locally advanced epithelioid sarcoma in patients not eligible for complete resection
Age Restrictions	16 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TECFIDERA**

### **Products Affected**

• TECFIDERA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., relapsing-remitting MS or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TEFLARO**

#### **Products Affected**

• TEFLARO

PA Criteria	Criteria Details
Exclusion Criteria	Known serious hypersensitivity to cephalosporin class
Required Medical Information	Diagnosis of one of the following A.) acute bacterial skin and skin structure infection and patient has documented culture and sensitivity to teflaro, or B.) community acquired pneumonia and patient has documented culture and sensitivity to teflaro
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	2 weeks
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TEGSEDI**

### **Products Affected**

• TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Platelet count less than 100,000 per microliter, B.) urinary protein to creatinine ratio (UPCR) of 1000 mg/g or higher
Required Medical Information	Diagnosis of Polyneuropathy of hereditary transthyretin-mediated amyloidosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TESTOSTERONE**

#### **Products Affected**

- *methyltestosterone oral*
- testosterone cypionate intramuscular solution 100 mg/ml, 200 mg/ml, 200 mg/ml (1 ml)
- testosterone enanthate intramuscular solution
- testosterone transdermal gel 10 mg/act (2%), 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)
- testosterone transdermal solution

	T
PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Carcinoma of the breast (males only) or prostate, B.) Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Hypogonadotropic hypogonadism, B.) Inoperable metastatic breast cancer in women who are postmenopausal, C.) Primary hypogonadism. Diagnosis of hypogonadism must be confirmed by a low-for-age serum testosterone (total or free) level defined by the normal laboratory reference value, or D.) Delayed puberty (testosterone enanthate)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135 PA20 C

Formulary ID:20153\_Version 12

## **TETRABENAZINE**

#### **Products Affected**

tetrabenazine

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Actively suicidal, B.) Untreated or inadequately treated depression, C.) Impaired hepatic function, D.) Concomitant use of monoamine oxidase inhibitors, E.) Concomitant use of reserpine or within 20 days of discontinuing reserpine
Required Medical Information	Diagnosis of chorea associated with Huntington's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **THALOMID**

#### **Products Affected**

• THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma that is newly diagnosed, or B.) Erythema nodosum leprosum (ENL)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TIBSOVO**

### **Products Affected**

• TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acute myeloid leukemia in relapsed or refractory patients, with susceptible isocitrate dehydrogenase-1 mutation, or B.) Acute myeloid leukemia in newly-diagnosed patients, with susceptible isocitrate dehydrogenase-1 mutation AND one of the following 1.) patient is 75 years or older, or 2.) patient has comorbidities that preclude intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TIGLUTIK**

#### **Products Affected**

• TIGLUTIK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of amyotrophic lateral sclerosis (ALS)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TOPICAL RETINOIDS**

#### **Products Affected**

- adapalene external cream
- adapalene external gel
- AVITA

- tretinoin external cream
- tretinoin external gel 0.01 %, 0.025 %

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate acne vulgaris
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TOREMIFENE**

#### **Products Affected**

• toremifene citrate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic breast cancer and patient must have previous inadequate response or intolerance to tamoxifen
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	6 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TRACLEER**

#### **Products Affected**

• TRACLEER ORAL TABLET SOLUBLE

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Receiving concomitant cyclosporine A or glyburide therapy, B.) Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal, or C.) Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND all of the following: A.) Patient has WHO Group I PAH, B.) Patient has New York Heart Association (NYHA) Functional Class II-IV, and C.) Female patients of reproductive potential must use two forms of reliable contraception
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TRELSTAR**

#### **Products Affected**

• TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer and used in palliative treatment
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# TRIKAFTA

## **Products Affected**

• TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis and patient has at least 1 F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene verified by an FDA-cleared CF mutation test
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TURALIO**

### **Products Affected**

• TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **TYKERB**

## **Products Affected**

• TYKERB

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) and One of the following A.) used in combination with capecitabine in a patient who has received prior therapy including an anthracycline, a taxane and trastuzumab, or B.) used in combination with letrozole in postmenopausal women for whom hormonal therapy is indicated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **TYMLOS**

### **Products Affected**

• TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of postmenopausal osteoporosis and one of the following A.) osteoporotic fracture or multiple risk factors for fracture, or B.) previous trial of/or contraindication to bisphosphonate
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: 12 months, max treatment 24 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **UPTRAVI**

### **Products Affected**

• UPTRAVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization and patient has tried and had an insufficient response to at least one other PAH agent (e.g., sildenafil) therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **VALCHLOR**

## **Products Affected**

VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (stage IA and IB mycosis fungoides-type) and patient has received prior skin-directed therapy (e.g. Topical corticosteroids, phototherapy, or topical nitrogen mustard)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **VARIZIG**

#### **Products Affected**

• VARIZIG INTRAMUSCULAR SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	History of hypersensitivity (including anaphylaxis or severe systemic reaction) to immune globulin or any component of the preparation
Required Medical Information	Diagnosis of post-exposure varicella (chickenpox) infection prophylaxis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **VENCLEXTA**

#### **Products Affected**

VENCLEXTA

#### • VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A inhibitor during the initial and titration phase in patients with CLL or SLL
Required Medical Information	Diagnosis of one of the following A.) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL), or B.) Newly-diagnosed acute myeloid leukemia (AML) and used in combination with azacitidine, decitabine or low-dose cytarabine in patients 75 years or older or who have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **VERZENIO**

### **Products Affected**

VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic, HER2-negative, hormone receptor-positive breast cancer AND One of the following: A.) For postmenopausal women must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy and patient has trial and failure or contraindication to Ibrance or Kisqali, B). For premenopausal or perimenopausal women must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy and patient has trial and failure or contraindication to Ibrance, C.) used as monotherapy for treatment of disease progression following endocrine therapy and patient has already received at least one prior chemotherapy regimen of Ibrance or Kisqali, D.) For postmenopausal women used as initial endocrine- based treatment in combination with an aromatase inhibitor and patient has trial and failure or contraindication to Kisqali or Ibrance, or E.) For premenopausal or perimenopausal women used as initial endocrine- based treatment in combination with an aromatase inhibitor and patient has trial and failure or contraindication to Kisqali
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

Y0135\_PA20\_C Formulary ID:20153\_Version 12 Last Updated: 05/19/2020

Effective Date: 06/01/2020

## **VIGABATRIN**

#### **Products Affected**

vigabatrin

#### VIGADRONE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Infantile spasms, or B.) Refractory complex partial seizures and the drug is being used as adjunctive therapy in patients who have responded inadequately to several alternative treatments
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **VITRAKVI**

#### **Products Affected**

VITRAKVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or surgically unresectable neurotrophic receptor tyrosine kinase (NTRK) gene fusion-positive solid tumors and used in patients with unsatisfactory alternative treatments or who have progressed following treatment
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **VIZIMPRO**

### **Products Affected**

• VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	First line treatment of metastatic non-small cell lung cancer with confirmed epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **VOTRIENT**

#### **Products Affected**

VOTRIENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) advanced renal cell carcinoma, or B.) advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., doxorubicin, dacarbazine, ifosfamide, epirubicin, docetaxel, or vinorelbine)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **VYNDAMAX**

### **Products Affected**

VYNDAMAX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of wild type or hereditary transthyretin related familial amyloid cardiomyopathy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **XALKORI**

### **Products Affected**

XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **XELJANZ**

#### **Products Affected**

XELJANZ

#### • XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) Moderate to severe rheumatoid arthritis (RA), B.) Active psoriatic arthritis, or C.) Moderate to severe ulcerative colitis (UC).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **XGEVA**

### **Products Affected**

• XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia (calcium less than 8.0 mg/dL)
Required Medical Information	Diagnosis of one of the following A.) bone metastases from a solid tumor, B.) giant cell tumor of the bone that is unresectable or where surgical resection is likely to result in severe morbidity, C.) hypercalcemia of malignancy refractory to bisphosphonate therapy, or D.) multiple myeloma used for the prevention of skeletal related events
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **XOLAIR**

#### **Products Affected**

XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic urticaria in patients who remain symptomatic despite H1 antihistamine therapy, or B.) Moderate to severe persistent asthma in patients with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms are inadequately controlled with inhaled corticosteroids
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, immunologist, pulmonologist, or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **XOSPATA**

### **Products Affected**

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia, with presence of FLT3 mutation as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **XPOVIO**

#### **Products Affected**

- XPOVIO (60 MG ONCE WEEKLY)
- XPOVIO (100 MG ONCE WEEKLY) XPOVIO (80 MG ONCE WEEKLY)
  - XPOVIO (80 MG TWICE WEEKLY)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory multiple myeloma being used in combination with dexamethasone in patient who has received at least 4 prior therapies and is refractory to at least 2 proteasome inhibitors, at least 2 immunomodulatory agents, and an anti-CD38 monoclonal antibody.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153 Version 12

## **XTANDI**

#### **Products Affected**

• XTANDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) castration-resistant prostate cancer (CRPC), or B) metastatic castration-sensitive prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **XYREM**

#### **Products Affected**

• XYREM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency
Required Medical Information	Diagnosis of cataplexy and excessive daytime sleepiness in patients with narcolepsy
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **YONSA**

#### **Products Affected**

• YONSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, or B.) Patients with severe baseline hepatic impairment (Child-Pugh Class C)
Required Medical Information	Diagnosis of metastatic castration-resistant prostate cancer and All of the following 1.) used in combination with methylprednisolone, and 2.) documented history of trial with, inadequate treatment response, adverse event, or contraindication to Zytiga (Abiraterone)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ZARXIO**

## **Products Affected**

• ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) congenital, cyclic, or idiopathic neutropenia, B) severe febrile neutropenia (FN) with the following: Has not received prophylactic pegfilgrastim and Used as adjunct to appropriate antibiotics in high-risk patients and any one of the following: 65 years or older, Uncontrolled primary disease, Pneumonia, Hypotension and multiorgan dysfunction (sepsis syndrome), Invasive fungal infection, Hospitalization when developed fever, Prior FN, Severe (ANC less than 100/mcL) or anticipated prolonged (more than 10 days) neutropenia, C) Autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, D) Undergoing myeloablative chemotherapy followed by autologous or allogeneic BMT, E) Acute myeloid leukemia and will be given after completion of induction or consolidation chemotherapy, F) Acute lymphoblastic leukemia and will be given after completion of the first few days of chemotherapy of the initial induction or first post-remission course, G) Myelodysplastic syndrome with severe neutropenia and recurrent infection, H) Receiving radiation therapy, not on chemotherapy, and expected to have prolonged delays in treatment due to neutropenia, I) Neutropenia associated with HIV infection and antiretroviral therapy, J) Aplastic anemia, K)Primary prophylaxis of FN in one of the following patients: 20% or higher risk of FN based on chemotherapy regimen OR Less than 20% risk of FN based on chemotherapy regimen with one of the following: 65 years or older, Poor performance status, Poor nutritional status, Previous FN, Extensive prior treatment including large radiation ports, Cytopenias due to bone marrow involvement by tumor, Administration of combined chemoradiotherapy, Presence of open wounds or active infections, Other serious comorbidities (including renal or liver dysfunction) or Receiving dose-dense chemotherapy regimen in breast or small cell lung cancer or non-Hodgkins lymphoma.

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

PA Criteria	Criteria Details
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ZEJULA**

#### **Products Affected**

ZEJULA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer and used for maintenance therapy in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin), or B.) advanced ovarian, fallopian tube, or primary peritoneal cancer and patient has been treated with 3 or more prior chemotherapy regimens, and cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA mutation, or genomic instability, and disease has progressed more than 6 months after response to the last platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or gynecologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ZELBORAF**

#### **Products Affected**

• ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test, or B.) Erdheim-Chester disease and patient has documented BRAF V600 mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ZOLINZA**

#### **Products Affected**

ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL) in patients who have progressive, persistent or recurrent disease on or following two systemic therapies (e.g., bexarotene, romidepsin, etc)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ZORTRESS**

#### **Products Affected**

ZORTRESS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Prevention of kidney transplant organ rejection, or B.) Prevention of liver transplant organ rejection
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescriber is experienced in immunosuppressive therapy and management of transplant patients.
Coverage Duration	12 months
Other Criteria	none
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

## **ZYDELIG**

#### **Products Affected**

ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic lymphocytic leukemia and all of the following: Used in combination with rituximab, patient has relapsed on at least one prior therapy (e.g., purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]), and patient does not have any co-morbidities that prevents the use of cytotoxic chemotherapy (i.e. severe neutropenia or thrombocytopenia, creatinine clearance less than 60 mL/minute), B.) Non-Hodgkins lymphoma (Follicular, B-Cell) and the patient has relapsed on at least two prior systemic therapies (e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]), or C.) Small lymphocytic lymphoma and the patient has relapsed on at least two prior systemic therapies(e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine])
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# ZYKADIA

#### **Products Affected**

• ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

# **ZYTIGA**

#### **Products Affected**

• abiraterone acetate

• ZYTIGA ORAL TABLET 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant metastatic prostate cancer and used in combination with prednisone, or B.) High risk, castration-sensitive metastatic prostate cancer and used in combination with prednisone
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	N/A
Off-Label Uses	N/A

Y0135\_PA20\_C

Formulary ID:20153\_Version 12

#### PART B VERSUS PART D

#### **Products Affected**

- ABELCET INTRAVENOUS SUSPENSION 5 MG/ML
- acetylcysteine inhalation solution 10 %, 20 %
- acyclovir sodium intravenous solution 50 mg/ml
- albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml
- AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG
- amikacin sulfate injection solution 500 mg/2ml
- AMINOSYN II INTRAVENOUS SOLUTION 10 %
- AMINOSYN-PF INTRAVENOUS SOLUTION 10 %, 7 %
- amphotericin b intravenous solution reconstituted 50 mg
- ampicillin-sulbactam sodium intravenous solution reconstituted 15 (10-5) gm
- aprepitant oral capsule 125 mg, 40 mg, 80
   & 125 mg, 80 mg
- AZACTAM INJECTION SOLUTION RECONSTITUTED 2 GM
- AZASAN ORAL TABLET 100 MG, 75 MG
- azathioprine oral tablet 50 mg
- azithromycin intravenous solution reconstituted 500 mg
- budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml
- calcitonin (salmon) nasal solution 200 unit/act
- calcitriol oral capsule 0.25 mcg, 0.5 mcg
- calcitriol oral solution 1 mcg/ml
- cefoxitin sodium intravenous solution reconstituted 1 gm, 2 gm
- ceftriaxone sodium injection solution reconstituted 1 gm, 2 gm, 250 mg, 500 mg

- cinacalcet hcl oral tablet 30 mg, 60 mg,
   90 mg
- ciprofloxacin in d5w intravenous solution 200 mg/100ml
- clindamycin phosphate injection solution 300 mg/2ml, 900 mg/6ml
- CLINIMIX E/DEXTROSE (2.75/5) INTRAVENOUS SOLUTION 2.75 %
- CLINIMIX E/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX E/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX E/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %
- CLINIMIX E/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %
- CLINIMIX/DEXTROSE (4.25/10)
   INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %
- CLINIMIX/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %
- CLINISOL SF INTRAVENOUS SOLUTION 15 %
- cromolyn sodium inhalation nebulization solution 20 mg/2ml
- cyclophosphamide oral capsule 25 mg, 50 mg
- cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg
- cyclosporine modified oral solution 100 mg/ml
- cyclosporine oral capsule 100 mg, 25 mg
- DEPO-PROVERA INTRAMUSCULAR SUSPENSION 400 MG/ML
- dextrose intravenous solution 10 %, 5 %
- dextrose-nacl intravenous solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.225 %, 5-0.45 %, 5-0.9 %

- diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml
- DOXY 100 INTRAVENOUS SOLUTION RECONSTITUTED 100 MG
- ENGERIX-B INJECTION SUSPENSION 10 MCG/0.5ML, 20 MCG/ML
- ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG
- ERYTHROCIN LACTOBIONATE INTRAVENOUS SOLUTION RECONSTITUTED 500 MG
- fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-%
- FREAMINE HBC INTRAVENOUS SOLUTION 6.9 %
- GENGRAF ORAL CAPSULE 100 MG, 25 MG
- GENGRAF ORAL SOLUTION 100 MG/ML
- granisetron hcl oral tablet 1 mg
- HEPATAMINE INTRAVENOUS SOLUTION 8 %
- imipenem-cilastatin intravenous solution reconstituted 250 mg, 500 mg
- IMOVAX RABIES INTRAMUSCULAR INJECTABLE 2.5 UNIT/ML
- INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %
- ipratropium bromide inhalation solution 0.02 %
- ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml
- ISOLYTE-P IN D5W INTRAVENOUS SOLUTION
- ISOLYTE-S INTRAVENOUS SOLUTION
- kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.9 meq/l-%-%, 30-5-0.45 meq/l-%-%, 40-5-0.9 meq/l-%-%
- *kcl-lactated ringers-d5w intravenous solution 20 meq/l*

- levalbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/0.5ml, 1.25 mg/3ml
- levofloxacin in d5w intravenous solution 500 mg/100ml, 750 mg/150ml
- levofloxacin intravenous solution 25 mg/ml
- meperidine hcl injection solution 100 mg/ml, 25 mg/ml, 50 mg/ml
- meropenem intravenous solution reconstituted 1 gm, 500 mg
- methotrexate oral tablet 2.5 mg
- methotrexate sodium (pf) injection solution 50 mg/2ml
- methotrexate sodium injection solution 50 mg/2ml
- metronidazole in nacl intravenous solution 500-0.79 mg/100ml-%
- moxifloxacin hcl in nacl intravenous solution 400 mg/250ml
- mycophenolate mofetil oral capsule 250 mg
- mycophenolate mofetil oral suspension reconstituted 200 mg/ml
- mycophenolate mofetil oral tablet 500 mg
- mycophenolate sodium oral tablet delayed release 180 mg, 360 mg
- nafcillin sodium injection solution reconstituted 1 gm, 2 gm
- nafcillin sodium intravenous solution reconstituted 10 gm
- NEPHRAMINE INTRAVENOUS SOLUTION 5.4 %
- NORMOSOL-M IN D5W INTRAVENOUS SOLUTION
- NORMOSOL-R IN D5W INTRAVENOUS SOLUTION
- NORMOSOL-R PH 7.4 INTRAVENOUS SOLUTION
- nutrilipid intravenous emulsion 20 %
- ondansetron hcl oral solution 4 mg/5ml
- ondansetron hcl oral tablet 24 mg, 4 mg, 8 mg
- ondansetron oral tablet dispersible 4 mg, 8 mg

- oxacillin sodium in dextrose intravenous solution 1 gm/50ml, 2 gm/50ml
- paricalcitol oral capsule 1 mcg, 2 mcg, 4 mcg
- penicillin g sodium injection solution reconstituted 5000000 unit
- pentamidine isethionate inhalation solution reconstituted 300 mg
- pentamidine isethionate injection solution reconstituted 300 mg
- piperacillin sod-tazobactam so intravenous solution reconstituted 2.25 (2-0.25) gm, 3.375 (3-0.375) gm, 4.5 (4-0.5) gm, 40.5 (36-4.5) gm
- PLASMA-LYTE 148 INTRAVENOUS SOLUTION
- PLASMA-LYTE A INTRAVENOUS SOLUTION
- PLENAMINE INTRAVENOUS SOLUTION 15 %
- potassium chloride in dextrose intravenous solution 20-5 meq/l-%, 40-5 meq/l-%
- potassium chloride in nacl intravenous solution 20-0.45 meq/l-%, 20-0.9 meq/l-%, 40-0.9 meq/l-%
- potassium chloride intravenous solution 10 meq/100ml, 2 meq/ml, 2 meq/ml (20 ml), 20 meq/100ml, 40 meq/100ml
- PREMASOL INTRAVENOUS SOLUTION 10 %
- PROCALAMINE INTRAVENOUS SOLUTION 3 %
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PROSOL INTRAVENOUS SOLUTION 20 %
- PULMOZYME INHALATION SOLUTION 1 MG/ML
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 10 MCG/ML (1ML SYRINGE), 40 MCG/ML, 5 MCG/0.5ML

- rifampin intravenous solution reconstituted 600 mg
- SANDIMMUNE ORAL SOLUTION 100 MG/ML
- sirolimus oral solution 1 mg/ml
- sirolimus oral tablet 0.5 mg, 1 mg, 2 mg
- sodium chloride intravenous solution 0.45 %, 0.9 %, 3 %, 5 %
- tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg
- TDVAX INTRAMUSCULAR SUSPENSION 2-2 LF/0.5ML
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU
- tigecycline intravenous solution reconstituted 50 mg
- TPN ELECTROLYTES INTRAVENOUS CONCENTRATE
- TPN ELECTROLYTES INTRAVENOUS SOLUTION
- TRAVASOL INTRAVENOUS SOLUTION 10 %
- TREXALL ORAL TABLET 10 MG, 15 MG, 5 MG, 7.5 MG
- TROPHAMINE INTRAVENOUS SOLUTION 10 %
- vancomycin hcl in dextrose intravenous solution 750-5 mg/150ml-%
- vancomycin hcl in nacl intravenous solution 1-0.9 gm/200ml-%, 500-0.9 mg/100ml-%
- vancomycin hcl intravenous solution reconstituted 1 gm, 250 mg, 500 mg, 5000 mg, 750 mg
- VARUBI (180 MG DOSE) ORAL TABLET THERAPY PACK 2 X 90 MG
- VARUBI ORAL TABLET 90 MG
- voriconazole intravenous solution reconstituted 200 mg
- XATMEP ORAL SOLUTION 2.5 MG/ML
- ZERBAXA INTRAVENOUS SOLUTION RECONSTITUTED 1.5 (1-0.5) GM

• ZOSYN INTRAVENOUS SOLUTION 2-0.25 GM/50ML, 3-0.375 GM/50ML

#### **Details**

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

## **Alphabetical Listing**

A	AURYXIA21
ABELCET INTRAVENOUS	AUSTEDO22
SUSPENSION 5 MG/ML349	AVITA 298, 299, 300
abiraterone acetate 347, 348	AVONEX PEN INTRAMUSCULAR
acetylcysteine inhalation solution 10 %, 20	AUTO-INJECTOR KIT 193, 194, 195,
%349	196
acitretin 1	AVONEX PREFILLED
ACTIMMUNE3	INTRAMUSCULAR PREFILLED
acyclovir sodium intravenous solution 50	SYRINGE KIT 193, 194, 195, 196
mg/ml 349	AYVAKIT23
adapalene external cream 298, 299, 300	AZACTAM INJECTION SOLUTION
adapalene external gel 298, 299, 300	RECONSTITUTED 2 GM 349
adefovir dipivoxil112, 113, 114	AZASAN ORAL TABLET 100 MG, 75
ADEMPAS 4	MG349
AFINITOR DISPERZ	azathioprine oral tablet 50 mg 349
AFINITOR ORAL TABLET 10 MG 6, 7	azithromycin intravenous solution
albuterol sulfate inhalation nebulization	reconstituted 500 mg 349
solution (2.5 mg/3ml) 0.083%, 0.63	В
mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml 349	BALVERSA24
ALECENSA9	BANZEL25
alosetron hcl10	BARACLUDE ORAL SOLUTION 112,
ALUNBRIG13	113, 114
AMBISOME INTRAVENOUS	BENLYSTA SUBCUTANEOUS 26, 27
SUSPENSION RECONSTITUTED 50	BETASERON SUBCUTANEOUS KIT 193,
MG349	194, 195, 196
ambrisentan14	bexarotene28, 29
amikacin sulfate injection solution 500	bosentan 30
mg/2ml 349	BOSULIF31
AMINOSYN II INTRAVENOUS	BRAFTOVI ORAL CAPSULE 75 MG 32,
SOLUTION 10 % 349	33
AMINOSYN-PF INTRAVENOUS	BRUKINSA 34
SOLUTION 10 %, 7 % 349	budesonide inhalation suspension 0.25
amphotericin b intravenous solution	mg/2ml, 0.5 mg/2ml, 1 mg/2ml 349
reconstituted 50 mg 349	C
ampicillin-sulbactam sodium intravenous	CABLIVI
solution reconstituted 15 (10-5) gm 349	CABOMETYX 36
APOKYN SUBCUTANEOUS SOLUTION	calcitonin (salmon) nasal solution 200
CARTRIDGE 17, 18	unit/act349
aprepitant oral capsule 125 mg, 40 mg, 80 &	calcitriol oral capsule 0.25 mcg, 0.5 mcg 349
125 mg, 80 mg 349	calcitriol oral solution 1 mcg/ml 349
ARCALYST 19	CALQUENCE38
ARIKAYCE20	CAPRELSA 39
armodafinil47	CARBAGLU40

CAYSTON41	CORLANOR
cefoxitin sodium intravenous solution	COSENTYX (300 MG DOSE) 54, 55
reconstituted 1 gm, 2 gm	COSENTYX SENSOREADY (300 MG) 54,
ceftriaxone sodium injection solution	55 COTELLIC
reconstituted 1 gm, 2 gm, 250 mg, 500	COTELLIC
mg	cromolyn sodium inhalation nebulization
CIMZIA PREFILLED	solution 20 mg/2ml
CIMZIA SUBCUTANEOUS KIT 2 X 200	cyclophosphamide oral capsule 25 mg, 50
MG	mg
cinacalcet hcl oral tablet 30 mg, 60 mg, 90	cyclosporine modified oral capsule 100 mg,
mg	25 mg, 50 mg
CINRYZE	cyclosporine modified oral solution 100
ciprofloxacin in d5w intravenous solution	mg/ml
200 mg/100ml	cyclosporine oral capsule 100 mg, 25 mg350
clindamycin phosphate injection solution	CYSTARAN57
300 mg/2ml, 900 mg/6ml	D
CLINIMIX E/DEXTROSE (2.75/5)	dalfampridine er
INTRAVENOUS SOLUTION 2.75 %350	DALIRESP
CLINIMIX E/DEXTROSE (4.25/10)	daptomycin
INTRAVENOUS SOLUTION 4.25 %350	DAURISMO
CLINIMIX E/DEXTROSE (4.25/5)	deferasirox oral tablet 360 mg, 90 mg 61, 62,
INTRAVENOUS SOLUTION 4.25 %350	63
CLINIMIX E/DEXTROSE (5/15)	deferasirox oral tablet soluble 61, 62, 63
INTRAVENOUS SOLUTION 5 % 350	DEPO-PROVERA INTRAMUSCULAR
CLINIMIX E/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 % 350	SUSPENSION 400 MG/ML
	dextrose intravenous solution 10 %, 5 % 350
CLINIMIX/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 %350	dextrose-nacl intravenous solution 10-0.2 %,
CLINIMIX/DEXTROSE (4.25/5)	10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.225 %, 5-0.45 %, 5-0.9 %
INTRAVENOUS SOLUTION 4.25 %350	diclofenac sodium transdermal gel 3 % 64,
CLINIMIX/DEXTROSE (5/15)	65
INTRAVENOUS SOLUTION 5 % 350	diphtheria-tetanus toxoids dt intramuscular
CLINIMIX/DEXTROSE (5/20)	suspension 25-5 lfu/0.5ml350
INTRAVENOUS SOLUTION 5 % 350	DOXY 100 INTRAVENOUS SOLUTION
CLINISOL SF INTRAVENOUS	RECONSTITUTED 100 MG
SOLUTION 15 %	dronabinol
clobazam	DUPIXENT
CLOVIQUE	E
COMETRIQ (100 MG DAILY DOSE) 49,	ELIGARD165, 166, 167, 168
50, 51	EMSAM
COMETRIQ (140 MG DAILY DOSE) 49,	ENBREL MINI
50, 51	ENBREL SUBCUTANEOUS SOLUTION
COMETRIQ (60 MG DAILY DOSE)49, 50,	PREFILLED SYRINGE 69, 70, 71, 72
51	ENBREL SUBCUTANEOUS SOLUTION
COPIKTRA52	RECONSTITUTED 69, 70, 71, 72

ENBREL SURECLICK SUBCUTANEOUS	FREAMINE HBC INTRAVENOUS
SOLUTION AUTO-INJECTOR 69, 70,	SOLUTION 6.9 % 351
71, 72	FYCOMPA 101
ENDARI 74	G
ENGERIX-B INJECTION SUSPENSION	GAMMAGARD INJECTION SOLUTION
10 MCG/0.5ML, 20 MCG/ML 350	2.5 GM/25ML 141, 142, 143, 144, 145,
entecavir	146
ENTRESTO 75	GAMMAGARD S/D LESS IGA 141, 142,
ENVARSUS XR ORAL TABLET	143, 144, 145, 146
EXTENDED RELEASE 24 HOUR 0.75	GAMMAKED INJECTION SOLUTION 1
MG, 1 MG, 4 MG 350	GM/10ML 141, 142, 143, 144, 145, 146
EPIDIOLEX76	GAMMAPLEX INTRAVENOUS
<b>EPOGEN INJECTION SOLUTION 10000</b>	SOLUTION 10 GM/100ML, 10
UNIT/ML, 2000 UNIT/ML, 20000	GM/200ML, 20 GM/200ML, 5
UNIT/ML, 3000 UNIT/ML, 4000	GM/50ML 141, 142, 143, 144, 145, 146
UNIT/ML 82, 83, 84	GAMUNEX-C INJECTION SOLUTION 1
ERIVEDGE77	GM/10ML 141, 142, 143, 144, 145, 146
ERLEADA78	GENGRAF ORAL CAPSULE 100 MG, 25
erlotinib hcl79	MG351
ERYTHROCIN LACTOBIONATE	GENGRAF ORAL SOLUTION 100
INTRAVENOUS SOLUTION	MG/ML351
RECONSTITUTED 500 MG351	GILENYA ORAL CAPSULE 0.5 MG 102,
ESBRIET ORAL CAPSULE 80, 81	103
ESDICIE I ORAL CAI SULE 60, 61	
ESBRIET ORAL TABLET 801 MG 80, 81	GILOTRIF 104
•	
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF       104         glatiramer acetate       105         GLEOSTINE ORAL CAPSULE 10 MG,       100 MG, 40 MG       107, 108         GOCOVRI       109
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF       104         glatiramer acetate       105         GLEOSTINE ORAL CAPSULE 10 MG,       100 MG, 40 MG       107, 108         GOCOVRI       109         granisetron hcl oral tablet 1 mg       351         H
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF       104         glatiramer acetate       105         GLEOSTINE ORAL CAPSULE 10 MG,       100 MG, 40 MG       107, 108         GOCOVRI       109         granisetron hcl oral tablet 1 mg       351         H       HEPATAMINE INTRAVENOUS
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF       104         glatiramer acetate       105         GLEOSTINE ORAL CAPSULE 10 MG,       100 MG, 40 MG       107, 108         GOCOVRI       109       109         granisetron hcl oral tablet 1 mg       351         H       HEPATAMINE INTRAVENOUS         SOLUTION 8 %       351
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF
ESBRIET ORAL TABLET 801 MG 80, 81 everolimus	GILOTRIF

HUMIRA SUBCUTANEOUS PREFILLED	kcl-lactated ringers-d5w intravenous
SYRINGE KIT 117, 118, 119, 120, 121,	solution 20 meq/l
122	KISQALI (200 MG DOSE) 151, 152, 153,
I	154
IBRANCE	KISQALI (400 MG DOSE) 151, 152, 153,
ICLUSIG125	154
IDHIFA126	KISQALI (600 MG DOSE) 151, 152, 153,
imatinib mesylate127, 128	154
IMBRUVICA129	KISQALI FEMARA (400 MG DOSE). 151,
imipenem-cilastatin intravenous solution	152, 153, 154
reconstituted 250 mg, 500 mg 351	KISQALI FEMARA (600 MG DOSE). 151,
IMOVAX RABIES INTRAMUSCULAR	152, 153, 154
INJECTABLE 2.5 UNIT/ML 351	KISQALI FEMARA(200 MG DOSE) 151,
INCRELEX	152, 153, 154
INLYTA133	KORLYM 155
INREBIC	KUVAN156
INTRALIPID INTRAVENOUS	L
EMULSION 20 %, 30 % 351	LENVIMA (10 MG DAILY DOSE) 157,
INTRAROSA	158, 159, 160, 161
INTRON A 136, 137	LENVIMA (12 MG DAILY DOSE) 157,
ipratropium bromide inhalation solution	158, 159, 160, 161
0.02 %	LENVIMA (14 MG DAILY DOSE) 157,
ipratropium-albuterol inhalation solution	158, 159, 160, 161
0.5-2.5 (3) mg/3ml	LENVIMA (18 MG DAILY DOSE) 157,
IRESSA	158, 159, 160, 161
ISOLYTE-P IN D5W INTRAVENOUS	LENVIMA (20 MG DAILY DOSE) 157,
SOLUTION351	158, 159, 160, 161
ISOLYTE-S INTRAVENOUS SOLUTION	LENVIMA (24 MG DAILY DOSE) 157,
351	158, 159, 160, 161
itraconazole oral	LENVIMA (4 MG DAILY DOSE) 157, 158,
J	159, 160, 161
JADENU ORAL TABLET 180 MG . 61, 62,	LENVIMA (8 MG DAILY DOSE) 157, 158,
63	159, 160, 161
JADENU SPRINKLE 61, 62, 63	LEUKINE INJECTION SOLUTION
JAKAFI147	RECONSTITUTED 163, 164
JYNARQUE ORAL TABLET THERAPY	leuprolide acetate injection 165, 166, 167,
PACK148, 149	168
K	levalbuterol hel inhalation nebulization
KALYDECO150	solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25
kel in dextrose-nacl intravenous solution 10-	mg/0.5ml, 1.25 mg/3ml351
5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%,	levofloxacin in d5w intravenous solution
20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%-	500 mg/100ml, 750 mg/150ml 351
%, 30-5-0.45 meq/l-%-%, 40-5-0.45	levofloxacin intravenous solution 25 mg/ml
meq/l-%-%, 40-5-0.9 meq/l-%-% 351	
	lidocaine external ointment 169, 170, 171

lidocaine external patch 5 %	mycophenolate mofetil oral capsule 250 mg
lidocaine-prilocaine external cream 169, 170, 171	mycophenolate mofetil oral tablet 500 mg
linezolid intravenous solution 600 mg/300ml	mycophenolate sodium oral tablet delayed release 180 mg, 360 mg
LONSURF 176	N
LORBRENA 177 LUPRON DEPOT (1-MONTH) 165, 166,	nafcillin sodium injection solution
167, 168	reconstituted 1 gm, 2 gm
LUPRON DEPOT (3-MONTH) 165, 166,	reconstituted 10 gm
167, 168 LUPRON DEPOT (4-MONTH) 165, 166,	NATPARA198 NEPHRAMINE INTRAVENOUS
167, 168	SOLUTION 5.4 %
LUPRON DEPOT (6-MONTH) 165, 166,	NERLYNX
167, 168 LYNPARZA ORAL TABLET 178, 179	NEXAVAR       200         NINLARO       201
M	NORMOSOL-M IN D5W INTRAVENOUS
MATULANE	SOLUTION352
MAVYRET 115, 116 MAYZENT 181	NORMOSOL-R IN D5W INTRAVENOUS SOLUTION352
MEKINIST	NORMOSOL-R PH 7.4 INTRAVENOUS
MEKTOVI 183	SOLUTION352
meperidine hel injection solution 100	NORTHERA
mg/ml, 25 mg/ml, 50 mg/ml 351 meropenem intravenous solution	NOXAFIL ORAL SUSPENSION 203, 204 NUBEQA
reconstituted 1 gm, 500 mg351	NUCALA 206
methotrexate oral tablet 2.5 mg 351	NUEDEXTA207
methotrexate sodium (pf) injection solution	nutrilipid intravenous emulsion 20 % 352
50 mg/2ml	O OCTAGAM INTRAVENOUS SOLUTION
mg/2ml	1 GM/20ML, 2 GM/20ML 141, 142, 143,
methoxsalen rapid 190, 191	144, 145, 146
methyltestosterone oral 287, 288, 289, 290, 291, 292	octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50
metronidazole in nacl intravenous solution 500-0.79 mg/100ml-%352	mcg/ml, 500 mcg/ml 208, 209, 210 ODOMZO
miglustat	OFEV
modafinil47	OMNITROPE 110, 111
moxifloxacin hel in nacl intravenous	ondansetron hel oral solution 4 mg/5ml 352
solution 400 mg/250ml 352	ondansetron hel oral tablet 24 mg, 4 mg, 8 mg

ondansetron oral tablet dispersible 4 mg, 8	PLENAMINE INTRAVENOUS
mg 352	SOLUTION 15 % 353
OPSUMIT212	POMALYST230
ORFADIN213	posaconazole231
ORKAMBI214	potassium chloride in dextrose intravenous
OSPHENA215	solution 20-5 meq/l-%, 40-5 meq/l-% 353
OTREXUP SUBCUTANEOUS	potassium chloride in nacl intravenous
SOLUTION AUTO-INJECTOR 10	solution 20-0.45 meq/l-%, 20-0.9 meq/l-
MG/0.4ML, 12.5 MG/0.4ML, 15	%, 40-0.9 meq/1-%
MG/0.4ML, 17.5 MG/0.4ML, 20	potassium chloride intravenous solution 10
MG/0.4ML, 22.5 MG/0.4ML, 25	meq/100ml, 2 meq/ml, 2 meq/ml (20 ml),
MG/0.4ML 184, 185, 186, 187, 188, 189	20 meq/100ml, 40 meq/100ml 353
oxacillin sodium in dextrose intravenous	PRALUENT SUBCUTANEOUS
solution 1 gm/50ml, 2 gm/50ml 352	SOLUTION AUTO-INJECTOR 219, 220,
oxandrolone oral216	221
P	PREMASOL INTRAVENOUS SOLUTION
PANRETIN218	10 %
paricalcitol oral capsule 1 mcg, 2 mcg, 4	PROCALAMINE INTRAVENOUS
mcg352	SOLUTION 3 %
PEGASYS PROCLICK SUBCUTANEOUS	PROGRAF ORAL PACKET 0.2 MG, 1 MG
SOLUTION 180 MCG/0.5ML 222, 223,	
224	PROLASTIN-C INTRAVENOUS
PEGASYS SUBCUTANEOUS SOLUTION	SOLUTION RECONSTITUTED 11, 12
	PROMACTA ORAL PACKET 12.5 MG
penicillamine oral tablet 225, 226	
penicillin g sodium injection solution	PROMACTA ORAL TABLET 232, 233
reconstituted 5000000 unit 352	PROSOL INTRAVENOUS SOLUTION 20
pentamidine isethionate inhalation solution	%353
reconstituted 300 mg352	PULMOZYME INHALATION
pentamidine isethionate injection solution	SOLUTION 1 MG/ML 353
reconstituted 300 mg 352	PURIXAN
piperacillin sod-tazobactam so intravenous	Q
solution reconstituted 2.25 (2-0.25) gm,	quinine sulfate oral236, 237
3.375 (3-0.375) gm, 4.5 (4-0.5) gm, 40.5	R
(36-4.5) gm352	RABAVERT INTRAMUSCULAR
PIQRAY (200 MG DAILY DOSE)227, 228,	SUSPENSION RECONSTITUTED 353
229	ranolazine er
PIQRAY (250 MG DAILY DOSE)227, 228,	RASUVO SUBCUTANEOUS SOLUTION
229	AUTO-INJECTOR 10 MG/0.2ML, 12.5
PIQRAY (300 MG DAILY DOSE)227, 228,	MG/0.25ML, 15 MG/0.3ML, 17.5
229	MG/0.35ML, 20 MG/0.4ML, 22.5
PLASMA-LYTE 148 INTRAVENOUS	MG/0.45ML, 25 MG/0.5ML, 30
SOLUTION	MG/0.6ML, 7.5 MG/0.15ML 184, 185,
PLASMA-LYTE A INTRAVENOUS	186, 187, 188, 189
SOLUTION	RAVICTI 239

RECOMBIVAX HB INJECTION	SYMLINPEN 120 SUBCUTANEOUS
SUSPENSION 10 MCG/ML, 10	SOLUTION PEN-INJECTOR 268, 269,
MCG/ML (1ML SYRINGE), 40	270
MCG/ML, 5 MCG/0.5ML 353	SYMLINPEN 60 SUBCUTANEOUS
REGRANEX240	SOLUTION PEN-INJECTOR 268, 269,
REPATHA 219, 220, 221	270
REPATHA PUSHTRONEX SYSTEM. 219,	SYMPAZAN45
220, 221	SYNAREL271
REPATHA SURECLICK 219, 220, 221	SYNRIBO272
RETACRIT 82, 83, 84	T
REVLIMID241	tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg
rifampin intravenous solution reconstituted	353
600 mg 353	TAFINLAR
riluzole	TAGRISSO276
RINVOQ243	TALZENNA
ROZLYTREK244	TARGRETIN EXTERNAL28, 29
RUBRACA 245, 246	TASIGNA
RUCONEST247	TAVALISSE279
RYDAPT248	tazarotene external 280, 281, 282
S	TAZORAC EXTERNAL CREAM 0.05 %
SAMSCA249	280, 281, 282
SANDIMMUNE ORAL SOLUTION 100	TAZORAC EXTERNAL GEL280, 281, 282
MG/ML353	TAZVERIK283
SIGNIFOR250	TDVAX INTRAMUSCULAR
sildenafil citrate oral tablet 20 mg 251, 252	SUSPENSION 2-2 LF/0.5ML 353
sirolimus oral solution 1 mg/ml	TECFIDERA284
sirolimus oral tablet 0.5 mg, 1 mg, 2 mg 353	TEFLARO285
SKYRIZI (150 MG DOSE)	TEGSEDI286
sodium chloride intravenous solution 0.45	TENIVAC INTRAMUSCULAR
%, 0.9 %, 3 %, 5 %	INJECTABLE 5-2 LFU353
sofosbuvir-velpatasvir115, 116	testosterone cypionate intramuscular
SOLTAMOX	solution 100 mg/ml, 200 mg/ml, 200
SOMATULINE DEPOT256	mg/ml (1 ml) 287, 288, 289, 290, 291, 292
SOMAVERT258	testosterone enanthate intramuscular
SPRYCEL259	solution 287, 288, 289, 290, 291, 292
STELARA SUBCUTANEOUS	testosterone transdermal gel 10 mg/act (2%),
SOLUTION 45 MG/0.5ML 260, 261, 262	12.5 mg/act (1%), 20.25 mg/1.25gm
STELARA SUBCUTANEOUS	(1.62%), 20.25 mg/act (1.62%), 25
SOLUTION PREFILLED SYRINGE 260,	mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%),
261, 262	50 mg/5gm (1%) 287, 288, 289, 290, 291,
STIVARGA	292
SUTENT	testosterone transdermal solution 287, 288,
SYLATRON SUBCUTANEOUS KIT 200	289, 290, 291, 292
MCG, 300 MCG265, 266	tetrabenazine
SYMDEKO	THALOMID
5 1 1 1 D L 1 5 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	111111111111111111111111111111111111111

TIBSOVO296	VEMLIDY 112, 113, 114
tigecycline intravenous solution	VENCLEXTA315, 316
reconstituted 50 mg353	VENCLEXTA STARTING PACK 315, 316
TIGLUTIK297	VERZENIO317, 318
tobramycin inhalation 131	vigabatrin
toremifene citrate 301	VIGADRONE319
TPN ELECTROLYTES INTRAVENOUS	VITRAKVI 321
CONCENTRATE 353	VIZIMPRO 322
TPN ELECTROLYTES INTRAVENOUS	voriconazole intravenous solution
SOLUTION353	reconstituted 200 mg354
TRACLEER ORAL TABLET SOLUBLE	VOSEVI115, 116
303, 304	VOTRIENT
TRAVASOL INTRAVENOUS SOLUTION	VYNDAMAX
10 % 353	X
TRELSTAR MIXJECT 305	XALKORI325
tretinoin external cream 298, 299, 300	XATMEP ORAL SOLUTION 2.5 MG/ML
tretinoin external gel 0.01 %, 0.025 % 298,	
299, 300	XELJANZ326
TREXALL ORAL TABLET 10 MG, 15	XELJANZ XR 326
MG, 5 MG, 7.5 MG 353	XGEVA328
trientine hcl	XOLAIR 329
TRIKAFTA	XOSPATA 330
TROPHAMINE INTRAVENOUS	XPOVIO (100 MG ONCE WEEKLY) 331,
SOLUTION 10 %	332, 333
TURALIO	XPOVIO (60 MG ONCE WEEKLY) 331,
TYKERB	332, 333
TYMLOS 310	XPOVIO (80 MG ONCE WEEKLY) 331,
U	332, 333
UPTRAVI311	XPOVIO (80 MG TWICE WEEKLY) 331,
V	332, 333
VALCHLOR312	XTANDI
vancomycin hcl in dextrose intravenous	XYREM335
solution 750-5 mg/150ml-% 354	Y
vancomycin hel in nacl intravenous solution	YONSA336
1-0.9 gm/200ml-%, 500-0.9 mg/100ml-%	${f Z}$
354	ZARXIO 337, 338
vancomycin hcl intravenous solution	ZEJULA
reconstituted 1 gm, 250 mg, 500 mg, 5000	ZELBORAF340
mg, 750 mg 354	ZERBAXA INTRAVENOUS SOLUTION
VARIZIG INTRAMUSCULAR	RECONSTITUTED 1.5 (1-0.5) GM 354
SOLUTION313, 314	ZOLINZA
VARUBI (180 MG DOSE) ORAL	ZORTRESS
TABLET THERAPY PACK 2 X 90 MG	ZOSYN INTRAVENOUS SOLUTION 2-
354	0.25 GM/50ML, 3-0.375 GM/50ML 354
VARUBI ORAL TABLET 90 MG 354	ZYDELIG

ZYKADIA ORAL TABLET ...... 345, 346

ZYTIGA ORAL TABLET 500 MG347, 348